DISPLAY COPY

# ZOOPLANKTON ABUNDANCE IN THE CENTRAL PACIFIC

BY JOSEPH E. KING AND JOAN DEMOND

FISHERY BULLETIN 82

UNITED STATES DEPARTMENT OF THE INTERIOR, Douglas McKay, *Secretary*

FISH AND WILDLIFE SERVICE, John L. Farley, *Director*

## ABSTRACT

Zooplankton abundance in the central Pacific was investigated on four cruises of the *Hugh M. Smith* in 1950 and 1951. Quantitative oblique hauls were made to 200 meters' depth, employing 1-meter nets of 30xxx grit gauze. Composition of the collections was remarkably uniform and averaged by number 57 percent Copepoda, 12 percent Chaetognatha, 6 percent Tunicata, 5 percent Euphausiacea, 4 percent Siphonophora, and 4 percent Foraminifera. An analysis of variance of zooplankton volumes demonstrated significant differences between day and night hauls, between cruises, and among latitudes but not between longitudes.

The greatest abundance, both by number and volume, of zooplankton occurred in the region of the Equator. The rich zone, extending from about 6° N. to 5° S. latitude, supported populations three to four times as great as more northerly or southerly latitudes. The greatest concentrations were found north of the Equator, when related to a "convergence;" when no marked convergence existed the peak of abundance was displaced a few degrees southward. The abundance of zooplankton was correlated with inorganic phosphate, oxygen, temperature, and thermocline depth. These environmental factors are influenced by upwelling associated with the equatorial divergence, which replenishes the supply of nutrients in the euphotic zone and creates favorable conditions for the growth of plant and animal life. While the data presented do not give a measure of the rate of production, they do provide a useful index to the relative productivity of different areas of the central Pacific.

UNITED STATES DEPARTMENT OF THE INTERIOR, Douglas McKay, *Secretary*
FISH AND WILDLIFE SERVICE, John L. Farley, *Director*

# ZOOPLANKTON ABUNDANCE IN THE CENTRAL PACIFIC

By Joseph E. King and Joan Demond

FISHERY BULLETIN 82

From Fishery Bulletin of the Fish and Wildlife Service

VOLUME 54

UNITED STATES GOVERNMENT PRINTING OFFICE • WASHINGTON : 1953

For sale by the Superintendent of Documents, U. S. Government Printing Office, Washington 25, D. C.
Price 25 cents.

## CONTENTS


| | Page |
|---|---|
| Collection of zooplankton samples | 112 |
| Areas sampled | 112 |
| Sampling in the field | 118 |
| Treatment of samples | 121 |
| Composition of the zooplankton | 122 |
| Zooplankton as food | 124 |
| Abundance of the zooplankton | 125 |
| Effects of the Equatorial Current system | 131 |
| Variation in zooplankton abundance | 132 |
| Effect of time of sampling | 133 |
| Variations between cruises, latitudes, and longitudes | 135 |
| Correlations with environmental factors | 137 |
| Comparison with zooplankton abundance of other regions | 138 |
| Productivity | 141 |
| Summary and conclusions | 142 |
| Literature cited | 142 |

# ZOOPLANKTON ABUNDANCE IN THE CENTRAL PACIFIC

By JOSEPH E. KING, *Fishery Research Biologist*, and
JOAN DEMOND, *Fishery Aid*

The Pacific Oceanic Fishery Investigations is authorized and directed to gather information which will ensure the maximum development and utilization of the high-seas fishery resources of the United States territories and island possessions in the tropical and subtropical Pacific. One project of fundamental importance in the research program concerns the relative productivity of different areas of the sea.

Productivity has been defined by Ivlev (1945) as the "capacity of a body of water to produce a given quantity of organic matter in some particular form." A direct measurement of the "rate of production" (Clarke 1946) would require that all processes by which organic matter is built up and destroyed be known and that the rates of these processes be determined. This is difficult to do and seldom has been done even for enclosed bodies of water. In mid-ocean the difficulties are vastly greater. We believe, however, that relative productivity, or productivity as defined by Ivlev, may be estimated indirectly by measuring the amounts of basic chemical nutrients in the water and the standing crops of plankton and fish. This report considers the quantity of zooplankton, 1 of the 2 main constituents of the total plankton crop, and its relation to certain physical and chemical factors in the central Pacific environment.

Zooplankton is essential food for much of the vertebrate fauna of the sea. It is utilized both directly and indirectly by tunas (the group of fish presently under study by these investigations). Kishinouye (1924) and Imamura (1949) have shown that zooplankton is prominent in the food of juvenile tunas. A variety of zooplankton organisms has also been observed in the food of adult tunas (Kishinouye 1917; Beebe 1936; Suyehiro 1942; Clemens and Wilby 1946; Reintjes and King 1953). The bulk of the zooplankton, however, reaches the tuna indirectly, being utilized by plankton-feeding animals which are in turn eaten by the tunas.

Potential food-fish resources are likely to exist in proportion to the amount of substance available for their nutriment. When vast areas of the sea are to be investigated, the several physical, chemical, and biotic properties of water associated with the production of nutriment for fish can be more readily and reliably surveyed than the abundance of the fish themselves. This report is concerned with the zooplankton from the particular viewpoint of its usefulness as an indicator of the relative productivity of the various portions of the area covered.

The literature includes a number of papers dealing with the plankton of the tropical and subtropical Pacific. One of the most valuable of these is the report by Graham (1941) on plankton collections taken by the *Carnegie* in the eastern and central Pacific. Kramer (1906) reported on a series of collections extending from Samoa to the Marshall Islands. Jesperson (1935) described results obtained by the *Dana* while traversing a series of stations reaching from Panama to the western Pacific, south of the Equator. For the western Pacific there are the publications of Matsuya (1937), Motoda (1940), Haneda (1942), and Tokioka (1942), which deal mainly with the plankton of lagoon, bay, and coastal waters but also provide some data on offshore plankton. The several papers of Marshall (1933), Russell (1934), and Russell and Colman (1931, 1934, 1935) supply a wealth of information on the plankton of the Great Barrier Reef Lagoon, but little on oceanic plankton. The papers of Johnson (1949) on the plankton of Bikini, and Sargent and Austin (1949) on the productivity of an atoll in the northern Marshalls, also deal primarily with the lagoon environment. The California Cooperative Sardine Research Program—a cooperative undertaking of the Scripps Institution of Oceanography, the United States Fish and Wild-

life Service, the California Department of Fish and Game, the California Academy of Sciences, and the Hopkins Marine Station of Stanford University—is collecting considerable information on the zooplankton of the eastern Pacific Ocean in subtropic and temperate latitudes. This organization has supplied our laboratory with copies of its unpublished data, which, when compared with our own results, show interesting differences between the zooplankton crops of the eastern and central Pacific. Reports of many of the expeditions which have entered the Pacific, such as those of the *Challenger*, the *Albatross*, and the *Meteor*, provide extensive information on the systematics and distribution of species or groups of the zooplankton, but supply little quantitative data that may be used to evaluate the zooplankton crop in the different regions visited.

The authors wish to acknowledge their indebtedness to Dr. Milner B. Schaefer,[1] under whose direction this project was initiated, and to Dr. Albert L. Tester[2] for his very valuable assistance in the statistical phases of the study and his constructive criticism of the manuscript.

We are also grateful to fellow staff members an the officers and crew of the *Hugh M. Smith* f their interest and efforts in obtaining this exte sive series of collections.

[1] Formerly chief, Section of Research and Development, Pacific Oceanic Fishery Investigations; presently director of Investigations, Inter-American Tropical Tuna Commission.

[2] Professor of Zoology, University of Hawaii.

## COLLECTION OF ZOOPLANKTON SAMPLES

### AREAS SAMPLED

This study is based on 210 collections made the central Pacific on cruises 2, 5, 7, and 8 of t United States Fish and Wildlife Service ves *Hugh M. Smith* in 1950 and 1951. Cruise 2 w made in January and February 1950; cruise 5 June, July, and August 1950; cruise 7 in Octob and November 1950; and cruise 8 in Januar February, and March, 1951. Thus, there were cruises (2 and 8) at a time corresponding to t northern winter season, and 1 cruise each (5 a 7) for summer and autumn. The approxima locations of the stations are shown in figures and 1b, and more exact positions are given tables 1, 2, 3, and 4. The data are distributed i long north-south sections, 6 of which cross t Equator, and in a number of shorter series stations (cruise 8).




FIGURE 1a.—Plankton-station positions of the *Hugh M. Smith*. A, cruise 2, January–March, 1950. B, cruise 5, June–August 1950.




FIGURE 1b.—Plankton-station positions of the *Hugh M. Smith*. C, cruise 7, October–November, 1950. D, cruise 8, January–March, 1951.

TABLE 1.—*Cruise 2: Estimated numbers and volumes of zooplankton collected, January to March, 1950*

| Station No. | Position | | Date | Time [1] | Water strained, in m.³ | Number of organisms per m.³ | Wet volume, cc. per m.³ |
|---|---|---|---|---|---|---|---|
| | Latitude | Longitude | | | | | |
| . Oblique tows to 200 m. (1-meter net, 30xxx mesh): | | | | | | | |
| 1 | 23°54′ N. | 166°51′ W. | Jan. 26 | 1100–1128 | 757.2 | 15 | 0.0079 |
| 3 | 20°59′ N. | 168°16′ W. | Jan. 27 | 1401–1419 | 539.2 | 20 | .0137 |
| 7 | 13°54′ N. | 171°36′ W. | Jan. 30 | 0326–0351 | 937.5 | 15 | .0115 |
| 9 | 12°02′ N. | 172°04′ W. | do | 2315–2337 | 837.5 | 15 | .0129 |
| 11 | 10°02′ N. | 171°55′ W. | Jan. 31 | 1946–2003 | 544.5 | 9 | .0092 |
| 13 | 8°00′ N. | 171°48′ W. | Feb. 1 | 1745–1805 | 631.2 | 25 | .0177 |
| 15 | 6°07′ N. | 171°56′ W. | Feb. 3 | 1300–1328 | 1,030.0 | 45 | .0285 |
| 17 | 3°58′ N. | 172°00′ W. | Feb. 4 | 0855–0921 | 837.5 | 43 | .0349 |
| 19 | 2°00′ N. | 172°05′ W. | Feb. 5 | 0245–0321 | 1,047.0 | 61 | .0714 |
| 21 | 0°00 | 171°59′ W. | Feb. 6 | 0250–0326 | 1,104.0 | 109 | .0826 |
| 23 | 2°00′ S. | 172°02′ W. | do | 2235–2257 | 782.5 | 25 | .0549 |
| 25 | 4°00′ S. | 171°58′ W. | Feb. 7 | 1646–1712 | 694.0 | 44 | .0236 |
| . Oblique tows to 200 m. (2-meter net, 3/16-inch mesh, rear section 18xxx mesh): | | | | | | | |
| 2 | 23°10′ N. | 167°00′ W. | Jan. 26 | 1921–1959 | 6,766.0 | 1 | .0003 |
| 6 | 15°00′ N. | 171°06′ W. | Jan. 29 | 1746–1811 | 3,852.4 | .8 | .0019 |
| 8 | 13°00′ N. | 172°00′ W. | Jan. 30 | 1213–1240 | 3,195.5 | .3 | .0009 |
| 10 | 11°03′ N. | 172°00′ W. | Jan. 31 | 1035–1056 | 3,610.5 | 1 | .0032 |
| 12 | 9°00′ N. | 172°00′ W. | Feb. 1 | 0546–0635 | 8,951.4 | 7 | .0040 |
| 14 | 7°00′ N. | 172°00′ W. | Feb. 2 | 1827–1858 | 3,716.6 | 3 | .0081 |
| 16 | 5°02′ N. | 172°01′ W. | Feb. 4 | 0034–0112 | 5,232.2 | 3 | .0101 |
| 18 | 2°56′ N. | 172°04′ W. | do | 1824–1855 | 4,644.9 | 10 | .0117 |
| 20 | 1°01′ N. | 172°09′ W. | Feb. 5 | 1624–1647 | 2,369.9 | 6 | .0122 |
| 22 | 1°01′ S. | 172°00′ W. | Feb. 6 | 1236–1259 | 2,453.4 | 5 | .0114 |
| 24 | 3°00′ S. | 172°04′ W. | Feb. 7 | 0755–0779 | 3,226.8 | 5 | .0146 |
| 26 | 5°02′ S. | 171°53′ W. | Feb. 8 | 0230–0252 | 2,579.6 | 1 | .0012 |
| . Surface tows (1-meter net, 30xxx mesh): | | | | | | | |
| 30 | 4°55′ S. | 158°13′ W. | Feb. 18 | 0545–0616 | 1,300.5 | 83 | .0451 |
| 32 | 3°06′ S. | 158°21′ W. | Feb. 19 | 0212–0243 | 1,067.0 | 38 | .0370 |
| 34 | 1°07′ S. | 158°28′ W. | do | 2200–2232 | 1,231.0 | 57 | .0911 |
| 36 | 1°02′ N. | 158°05′ W. | Feb. 20 | 1030–1101 | 1,357.2 | 64 | .0721 |
| 38 | 3°02′ N. | 157°51′ W. | Feb. 22 | 1820–1850 | 1,035.5 | 58 | .0759 |
| 40 | 5°00′ N. | 157°59′ W. | Feb. 23 | 1255–1326 | 1,139.0 | 26 | .0088 |
| 42 | 7°00′ N. | 157°58′ W. | Feb. 24 | 0725–0756 | 1,148.0 | 27 | .0082 |
| 44 | 8°53′ N. | 157°54′ W. | Feb. 25 | 0025–0056 | 1,112.0 | 62 | .0622 |
| 46 | 11°02′ N. | 158°06′ W. | do | 2220–2302 | 1,284.8 | 19 | .0139 |
| 48 | 13°02′ N. | 157°51′ W. | Feb. 26 | 1830–1901 | 1,084.5 | 16 | .0037 |
| 50 | 14°57′ N. | 157°59′ W. | Feb. 27 | 1325–1356 | 1,332.5 | 5 | .0022 |
| 52 | 19°00′ N. | 157°42′ W. | Mar. 1 | 0215–0246 | 1,310.0 | 23 | .0176 |
| . Surface tows (1-meter net, 18xxx mesh): | | | | | | | |
| 31 | 3°53′ S. | 157°47′ W. | Feb. 18 | 1604–1636 | 1,066.0 | 17 | .0255 |
| 33 | 2°03′ S. | 158°28′ W. | Feb. 19 | 1231–1304 | 1,105.5 | 3 | .0054 |
| 35 | 0°04′ S. | 158°22′ W. | Feb. 20 | 0900–0931 | 1,307.8 | 4 | .0044 |
| 37 | 2°03′ N. | 158°08′ W. | Feb. 21 | 2045–2116 | 1,272.5 | 18 | .0520 |
| 39 | 4°03′ N. | 157°53′ W. | Feb. 23 | 0412–0442 | 1,205.5 | 24 | .0499 |
| 41 | 6°00′ N. | 158°03′ W. | do | 2150–2251 | 1,550.0 | 12 | .0329 |
| 43 | 8°02′ N. | 157°49′ W. | Feb. 24 | 1607–1638 | 1,190.5 | 6 | .0049 |
| 45 | 9°55′ N. | 157°54′ W. | Feb. 25 | 0952–1023 | 1,567.0 | 1 | .0003 |
| 47 | 12°01′ N. | 158°03′ W. | Feb. 26 | 0725–0755 | 1,772.8 | 1 | .0003 |
| 49 | 14°00′ N. | 157°51′ W. | Feb. 27 | 0427–0500 | 1,232.8 | 11 | .0190 |
| 51 | 17°01′ N. | 157°56′ W. | Feb. 28 | 0730–0801 | 1,620.0 | 2 | .0019 |
| 53 | 20°54′ N. | 158°04′ W. | Mar. 1 | 2056–2136 | 2,012.8 | 3 | .0053 |

[1] Time corresponding to +11 zone time (Greenwich Civil Time −11 hours) was used for all stations 1 to 26, and +10 zone time on stations 30 to 53.

TABLE 2.—*Cruise 5: Estimated numbers and volumes of zooplankton collected, June to August, 1950*

[All tows oblique, surface to 200 m. to surface; all nets with body of 30xxx grit gauze, cod end of 56xxx grit gauze]

| Station No. | Position | | Date | Time [1] | Water strained in m.³ | Number of organisms per m.³ | Wet volume, cc. per m.³ |
|---|---|---|---|---|---|---|---|
| | Latitude | Longitude | | | | | |
| | 27°00′ N. | 175°11′ W. | June 30 | 0945–1012 | 1,248.7 | 26 | 0.0118 |
| | 25°00′ N. | 174°10′ W. | July 1 | 0258–0325 | 1,322.7 | 46 | .0271 |
| | 22°58′ N. | 173°00′ W. | do | 2315–2337 | 1,343.9 | 40 | .0175 |
| | 21°00′ N. | 172°00′ W. | July 2 | 0415–0431 | 739.5 | 33 | .0254 |
| | 19°00′ N. | 171°52′ W. | July 3 | 0807–0831 | 1,254.8 | 35 | .0174 |
| | 17°01′ N. | 171°46′ W. | do | 1219–1246 | 1,434.4 | 22 | .0388 |
| | 15°00′ N. | 171°54′ W. | July 4 | 1649–1710 | 888.1 | 45 | .0216 |
| | 14°00′ N. | 171°57′ W. | July 5 | 0123–0145 | 854.8 | 40 | .0234 |
| | 13°00′ N. | 172°01′ W. | do | 0950–1019 | 1,327.5 | 32 | .0125 |
| | 12°00′ N. | 172°00′ W. | do | 1815–1833 | 882.8 | 14 | .0084 |
| | 11°03′ N. | 172°00′ W. | July 6 | 0315–0342 | 1,333.3 | 34 | .0374 |
| | 9°54′ N. | 172°02′ W. | do | 1213–1244 | 1,777.0 | 20 | .0252 |
| | 8°54′ N. | 172°00′ W. | do | 2012–2036 | 1,190.2 | 31 | .0301 |
| | 7°59′ N. | 171°57′ W. | July 7 | 0350–0420 | 1,623.7 | 18 | .0283 |
| | 6°59′ N. | 171°46′ W. | do | 1226–1258 | 1,579.4 | 27 | .0204 |
| | 6°02′ N. | 171°56′ W. | do | 2130–2201 | 1,382.2 | 30 | .0338 |
| | 5°00′ N. | 172°02′ W. | July 8 | 0845–0918 | 1,821.1 | 40 | .0226 |

Footnote at end of table.

TABLE 2.—*Cruise 5: Estimated numbers and volumes of zooplankton collected, June to August, 1950*—Con.

| Station | Position | | Date | Time [1] | Water strained, in m.³ | Number of organisms per m.³ | Wet volume cc. per m.³ |
|---|---|---|---|---|---|---|---|
| | Latitude | Longitude | | | | | |
| 18 | 4°00′ N. | 172°03′ W. | ---do---- | 1659–1729 | 1,346.2 | 65 | .03 |
| 19 | 3°00′ N. | 172°01′ W. | July 9 | 0036–0101 | 1,063.8 | 59 | .04 |
| 20 | 2°00′ N. | 171°53′ W. | ---do---- | 0838–0909 | 1,809.2 | 58 | .03 |
| 21 | 0°54′ N. | 172°12′ W. | ---do---- | 1746–1815 | 1,586.0 | 67 | .05 |
| 22 | 0°08′ S. | 172°02′ W. | July 10 | 0143–0214 | 1,943.8 | 94 | .06 |
| 23 | 1°00′ S. | 171°57′ W. | ---do---- | 0925–0951 | 1,256.9 | 107 | .07 |
| 24 | 2°00′ S. | 171°55′ W. | ---do---- | 1740–1809 | 1,406.4 | 52 | .04 |
| 25 | 2°59′ S. | 171°59′ W. | July 11 | 0133–0201 | 1,309.9 | 68 | .06 |
| 26 | 4°05′ S. | 171°56′ W. | ---do---- | 1055–1127 | 1,585.3 | 36 | .02 |
| 27 | 5°04′ S. | 171°58′ W. | ---do---- | 2110–2141 | 1,632.8 | 27 | .03 |
| 28 | 4°56′ S. | 158°02′ W. | July 28 | 0556–0614 | 1,384.8 | 20 | .00 |
| 29 | 4°00′ S. | 158°03′ W. | ---do---- | 1324–1338 | 642.8 | 47 | .02 |
| 30 | 3°03′ S. | 158°00′ W. | ---do---- | 2137–2200 | 873.7 | 46 | .04 |
| 31 | 2°03′ S. | 158°00′ W. | July 29 | 0550–0607 | 648.3 | 45 | .02 |
| 32 | 1°00′ S. | 158°01′ W. | ---do---- | 1418–1436 | 996.0 | 43 | .02 |
| 33 | 0°02′ S. | 157°58′ W. | ---do---- | 2221–2239 | 773.0 | 76 | .08 |
| 34 | 0°57′ N. | 157°57′ W. | July 30 | 0646–0715 | 1,528.5 | 44 | .03 |
| 35 | 2°00′ N. | 158°07′ W. | ---do---- | 1532–1553 | 851.2 | 62 | .06 |
| 36 | 3°00′ N. | 157°58′ W. | ---do---- | 2302–2324 | 972.8 | 93 | .07 |
| 37 | 4°02′ N. | 158°03′ W. | July 31 | 0640–0707 | 1,652.5 | 76 | .07 |
| 38 | 5°00′ N. | 158°00′ W. | ---do---- | 1750–1816 | 1,175.4 | 74 | .06 |
| 39 | 6°00′ N. | 157°57′ W. | Aug. 1 | 0144–0214 | 1,313.7 | 35 | .04 |
| 40 | 7°00′ N. | 157°57′ W. | ---do---- | 0854–0919 | 954.3 | 43 | .02 |
| 41 | 8°00′ N. | 157°50′ W. | ---do---- | 1620–1704 | 3,361.5 | 31 | .02 |
| 42 | 9°00′ N. | 157°55′ W. | Aug. 2 | 0020–0046 | 1,136.6 | 38 | .04 |
| 43 | 10°00′ N. | 157°53′ W. | ---do---- | 0744–0812 | 1,597.8 | 25 | .02 |
| 44 | 11°00′ N. | 157°55′ W. | ---do---- | 1546–1615 | 1,141.7 | 40 | .02 |
| 45 | 11°59′ N. | 158°04′ W. | ---do---- | 2343–0008 | 1,034.5 | 32 | .03 |
| 46 | 13°00′ N. | 157°58′ W. | Aug. 3 | 0800–0827 | 1,455.8 | 19 | .01 |
| 47 | 14°00′ N. | 157°54′ W. | ---do---- | 1639–1709 | 1,305.7 | 18 | .02 |
| 48 | 14°58′ N. | 157°55′ W. | Aug. 4 | 0253–0311 | 592.8 | 19 | .02 |
| 49 | 17°00′ N. | 158°08′ W. | ---do---- | 1741–1805 | 934.8 | 38 | .02 |
| 50 | 19°02′ N. | 157°59′ W. | Aug. 5 | 1006–1031 | 1,035.0 | 29 | .03 |
| 51 | 20°53′ N. | 158°02′ W. | ---do---- | 2300–2324 | 852.2 | 37 | .02 |

[1] +11 zone time on stations 1 to 27, +10 zone time on stations 28 to 51.

TABLE 3.—*Cruise 7: Estimated numbers and volumes of zooplanton collected, October to November 1950*

[All tows oblique, surface to 200 m. to surface; all nets with body of 30xxx grit gauze, cod end of 56xxx grit gauze]

| Station No. | Position | | Date | Time [1] | Water strained, in m.³ | Number of organisms per m.³ | Wet volume cc. per m.³ |
|---|---|---|---|---|---|---|---|
| | Latitude | Longitude | | | | | |
| 1 | 20°30′ N. | 158°00′ W. | Oct. 17 | 1925–1955 | 952.8 | 44 | 0.03 |
| 2 | 19°10′ N. | 158°02′ W. | Oct. 18 | 0532–0600 | 897.3 | 44 | .16 |
| 3 | 18°13′ N. | 158°00′ W. | ---do---- | 1326–1355 | 1,112.5 | 30 | .09 |
| 4 | 17°10′ N. | 158°00′ W. | ---do---- | 2159–2225 | 683.7 | 27 | .01 |
| 5 | 16°11′ N. | 157°57′ W. | Oct. 19 | 0506–0530 | 699.0 | 21 | .00 |
| 6 | 15°02′ N. | 157°58′ W. | ---do---- | 1250–1318 | 999.2 | 14 | .00 |
| 7 | 14°03′ N. | 157°59′ W. | ---do---- | 1921–1955 | 1,220.6 | 27 | .00 |
| 8 | 12°57′ N. | 157°57′ W. | Oct. 20 | 0330–0353 | 730.6 | 64 | .00 |
| 9 | 12°25′ N. | 158°00′ W. | ---do---- | 1729–1804 | 1,217.9 | 38 | .00 |
| 10 | 11°08′ N. | 158°01′ W. | Oct. 21 | 1526–1556 | 1,373.5 | 42 | .00 |
| 11 | 10°01′ N. | 157°55′ W. | Oct. 22 | 0724–0804 | 1,426.0 | 40 | .00 |
| 12 | 8°52′ N. | 157°45′ W. | Oct. 23 | 0722–0751 | 956.9 | 37 | .00 |
| 13 | 8°05′ N. | 157°15′ W. | Oct. 24 | 0742–0813 | 1,413.5 | 19 | .00 |
| 14 | 7°17′ N. | 157°04′ W. | Oct. 25 | 0727–0805 | 1,698.8 | 32 | .00 |
| 15 | 5°57′ N. | 157°05′ W. | Oct. 26 | 0737–0810 | 1,572.1 | 59 | .00 |
| 16 | 5°19′ N. | 157°20′ W. | Oct. 27 | 0725–0809 | 2,206.2 | 65 | .00 |
| 17 | 4°23′ N. | 157°30′ W. | Oct. 28 | 0912–0945 | 1,872.1 | 40 | .00 |
| 18 | 3°17′ N. | 157°53′ W. | Oct. 29 | 0802–0837 | 1,314.0 | 50 | .00 |
| 19 | 2°01′ N. | 157°34′ W. | Nov. 1 | 0812–0844 | 1,587.3 | 75 | .00 |
| 20 | 6°07′ N. | 162°06′ W. | Nov. 6 | 0950–1028 | 1,930.3 | 28 | .00 |
| 21 | 6°13′ N. | 163°05′ W. | Nov. 7 | 0810–0837 | 1,565.8 | 63 | .00 |
| 22 | 6°59′ N. | 163°54′ W. | Nov. 8 | 0808–0841 | 1,749.1 | 22 | .00 |
| 23 | 7°24′ N. | 164°25′ W. | Nov. 9 | 0806–0838 | 1,442.4 | 27 | .00 |
| 24 | 6°31′ N. | 165°45′ W. | Nov. 10 | 0813–0847 | 1,521.5 | 23 | .00 |

+10 zone time on stations 1 to 21, +11 zone time on stations 22 to 24.

TABLE 4.—*Cruise 8: Estimated numbers and volumes of zooplankton collected, January to March, 1951*

[All tows oblique, surface to 200 m. to surface; all nets with body of 30xxx grit gauze, cod end of 56xxx grit gauze]

| Station No. | Position | | Date | Time [1] | Water strained, in m.³ | Number of organisms per m.³ | Wet volume, cc. per m.³ |
|---|---|---|---|---|---|---|---|
| | Latitude | Longitude | | | | | |
| | 20°48′ N. | 157°30′ W. | Jan. 14 | 2012–2048 | 937.3 | 27 | 0.0154 |
| | 18°47′ N. | 158°01′ W. | Jan. 15 | 1449–1504 | 625.9 | 23 | .0105 |
| | 14°30′ N. | 158°00′ W. | Jan. 17 | 0925–0940 | 433.1 | 43 | .0236 |
| | 12°57′ N. | 157°58′ W. | do | 2251–2318 | 819.0 | 46 | .0310 |
| | 11°59′ N. | 157°50′ W. | Jan. 18 | 0811–0836 | 1,363.0 | 24 | .0094 |
| | 11°00′ N. | 157°53′ W. | do | 1633–1658 | 1,399.9 | 36 | .0093 |
| | 10°00′ N. | 157°56′ W. | Jan. 19 | 0224–0255 | 1,830.5 | 30 | .0251 |
| | 9°00′ N. | 157°58′ W. | do | 1154–1216 | 985.1 | 33 | .0166 |
| | 8°04′ N. | 157°58′ W. | do | 2024–2047 | 1,244.0 | 31 | .0190 |
| | 7°08′ N. | 157°58′ W. | Jan. 20 | 0615–0645 | 1,299.2 | 74 | .0477 |
| | 5°59′ N. | 158°00′ W. | do | 1505–1537 | 1,807.5 | 34 | .0186 |
| | 5°00′ N. | 158°00′ W. | do | 2338–2357 | 975.3 | 32 | .0174 |
| | 3°55′ N. | 157°54′ W. | Jan. 21 | 1100–1118 | 984.1 | 31 | .0118 |
| | 3°02′ N. | 157°57′ W. | do | 2135–2155 | 1,018.2 | 43 | .0255 |
| | 2°00′ N. | 158°01′ W. | Jan. 22 | 0802–0831 | 1,625.2 | 30 | .0146 |
| | 0°58′ N. | 158°07′ W. | do | 1730–1754 | 1,529.1 | 33 | .0220 |
| | 0°01′ N. | 158°02′ W. | Jan. 23 | 0323–0356 | 1,925.6 | 32 | .0370 |
| | 0°55′ S. | 157°54′ W. | do | 1208–1233 | 1,509.4 | 90 | .0478 |
| | 2°00′ S. | 158°00′ W. | do | 2312–2336 | 1,273.4 | 80 | .0558 |
| | 3°04′ S. | 158°05′ W. | Jan. 24 | 0744–0808 | 1,270.9 | 48 | .0257 |
| | 4°00′ S. | 158°00′ W. | do | 1550–1614 | 1,070.4 | 36 | .0178 |
| | 5°00′ S. | 158°00′ W. | do | 2330–2350 | 885.8 | 46 | .0666 |
| | 6°01′ S. | 158°06′ W. | Jan. 25 | 0741–0806 | 1,234.2 | 48 | .0266 |
| | 7°00′ S. | 158°01′ W. | do | 1548–1616 | 1,352.6 | 23 | .0092 |
| | 6°57′ N. | 154°56′ W. | Jan. 31 | 0208–0238 | 1,521.4 | 39 | .0251 |
| | 5°58′ N. | 155°08′ W. | do | 1040–1060 | 948.4 | 35 | .0325 |
| | 5°00′ N. | 155°28′ W. | do | 1836–1902 | 1,483.2 | 35 | .0284 |
| | 4°04′ N. | 155°46′ W. | Feb. 1 | 0223–0250 | 1,254.0 | 49 | .0340 |
| | 3°06′ N. | 156°09′ W. | do | 1040–1061 | 1,676.0 | 66 | .0348 |
| | 2°00′ N. | 156°30′ W. | do | 1835–1902 | 1,556.8 | 46 | .0315 |
| | 6°50′ N. | 157°33′ W. | Feb. 3 | 0612–0639 | 1,680.3 | 22 | .0145 |
| | 6°04′ N. | 157°53′ W. | do | 1337–1404 | 1,401.1 | 21 | .0126 |
| | 5°00′ N. | 158°34′ W. | do | 2245–2309 | 1,245.7 | 31 | .0204 |
| | 4°05′ N. | 158°53′ W. | Feb. 4 | 0706–0727 | 1,063.9 | 41 | .0177 |
| | 3°03′ N. | 159°10′ W. | Feb. 6 | 0441–0513 | 1,881.2 | 21 | .0254 |
| | 1°57′ N. | 159°39′ W. | do | 1343–1419 | 1,619.8 | 26 | .0106 |
| | 6°59′ N. | 161°12′ W. | Feb. 7 | 1025–1049 | 1,744.5 | 29 | .0224 |
| | 5°55′ N. | 161°28′ W. | Feb. 8 | 2030–2056 | 1,914.2 | 33 | .0313 |
| | 4°52′ N. | 161°48′ W. | Feb. 9 | 0443–0513 | 1,835.9 | 33 | .0276 |
| | 3°58′ N. | 162°02′ W. | do | 1119–1143 | 1,488.0 | 17 | .0118 |
| | 2°58′ N. | 162°25′ W. | do | 1919–1946 | 1,819.0 | 30 | .0303 |
| | 2°00′ N. | 162°57′ W. | Feb. 10 | 0259–0327 | 1,514.6 | 31 | .0354 |
| | 6°44′ N. | 165°23′ W. | Feb. 11 | 1652–1718 | 1,330.2 | 21 | .0170 |
| | 5°51′ N. | 165°28′ W. | Feb. 12 | 0021–0048 | 1,647.2 | 31 | .0368 |
| | 4°58′ N. | 165°35′ W. | do | 0758–0824 | 1,432.3 | 23 | .0176 |
| | 3°59′ N. | 166°01′ W. | do | 1626–1653 | 1,743.6 | 22 | .0118 |
| | 3°00′ N. | 166°36′ W. | Feb. 13 | 0100–0130 | 1,760.7 | 37 | .0359 |
| | 2°04′ N. | 166°47′ W. | do | 0848–0913 | 1,618.2 | 26 | .0248 |
| | 1°02′ N. | 167°02′ W. | do | 1815–1839 | 1,116.2 | 52 | .0408 |
| | 0°01′ N. | 167°23′ W. | Feb. 14 | 0240–0308 | 1,433.5 | 47 | .0473 |
| | 1°05′ S. | 167°36′ W. | do | 1132–1156 | 1,420.4 | 23 | .0172 |
| | 2°01′ S. | 167°42′ W. | do | 1903–1930 | 1,785.5 | 37 | .0320 |
| | 3°05′ S. | 167°50′ W. | Feb. 15 | 0323–0353 | 1,740.7 | 34 | .0295 |
| | 4°00′ S. | 167°58′ W. | do | 1041–1109 | 2,008.7 | 20 | .0175 |
| | 5°03′ S. | 168°09′ W. | do | 1819–1848 | 1,942.1 | 19 | .0175 |
| | 5°59′ S. | 168°29′ W. | Feb. 16 | 0208–0241 | 2,424.3 | 33 | .0332 |
| | 7°04′ S. | 168°48′ W. | do | 0923–0953 | 2,308.6 | 24 | .0100 |
| | 14°30′ S. | 171°51′ W. | Feb. 26 | 1931–1958 | 1,876.1 | 28 | .0154 |
| | 11°56′ S. | 171°59′ W. | Feb. 27 | 1129–1157 | 1,918.4 | 26 | .0115 |
| | 10°05′ S. | 171°52′ W. | Feb. 28 | 0245–0309 | 1,735.6 | 41 | .0226 |
| | 8°00′ S. | 171°56′ W. | do | 1821–1846 | 1,452.5 | 29 | .0153 |
| | 7°01′ S. | 171°54′ W. | Mar. 1 | 0238–0303 | 1,045.2 | 52 | .0291 |
| | 5°59′ S. | 172°00′ W. | do | 1118–1144 | 1,399.9 | 22 | .0100 |
| | 5°00′ S. | 172°15′ W. | do | 1930–1955 | 1,507.6 | 22 | .0202 |
| | 4°00′ S. | 172°06′ W. | Mar. 2 | 0735–0758 | 1,510.7 | 18 | .0101 |
| | 3°00′ S. | 171°50′ W. | do | 1537–1605 | 1,510.7 | 20 | .0146 |
| | 2°00′ S. | 172°02′ W. | Mar. 3 | 1612–1641 | 1,593.0 | 21 | .0109 |
| | 0°57′ S. | 172°00′ W. | Mar. 4 | 0047–0115 | 1,392.3 | 43 | .0394 |
| | 0°09′ N. | 171°58′ W. | do | 0926–0947 | 1,066.0 | 56 | .0345 |
| | 1°04′ N. | 172°00′ W. | do | 1702–1727 | 979.6 | 73 | .0398 |
| | 1°54′ N. | 172°02′ W. | Mar. 5 | 0034–0101 | 1,854.7 | 57 | .0548 |
| | 2°48′ N. | 172°04′ W. | do | 0834–0858 | 1,332.4 | 48 | .0435 |
| | 3°57′ N. | 172°02′ W. | do | 1816–1842 | 1,221.9 | 44 | .0367 |
| | 4°56′ N. | 171°54′ W. | Mar. 6 | 0249–0315 | 1,621.0 | 51 | .0369 |
| | 5°55′ N. | 171°50′ W. | do | 1105–1130 | 1,428.2 | 43 | .0242 |
| | 6°58′ N. | 171°49′ W. | do | 1939–2011 | 1,907.6 | 24 | .0244 |
| | 7°56′ N. | 171°51′ W. | Mar. 7 | 0407–0433 | 1,498.1 | 22 | .0139 |
| | 9°00′ N. | 171°58′ W. | do | 1315–1339 | 1,539.7 | 13 | .0094 |
| | 10°02′ N. | 171°56′ W. | do | 2147–2213 | 1,538.1 | 22 | .0127 |
| | 11°02′ N. | 171°52′ W. | Mar. 8 | 0602–0629 | 1,576.9 | 23 | .0115 |
| 0 | 12°00′ N. | 171°52′ W. | do | 1417–1443 | 1,507.0 | 11 | .0097 |
| 1 | 13°01′ N. | 171°48′ W. | do | 2236–2301 | 1,720.5 | 20 | .0192 |
| 2 | 13°58′ N. | 171°24′ W. | Mar. 9 | 0650–0713 | 1,274.5 | 15 | .0096 |
| 3 | 15°00′ N. | 170°52′ W. | do | 1557–1619 | 1,313.9 | 23 | .0100 |
| 4 | 16°58′ N. | 169°42′ W. | Mar. 10 | 0913–0940 | 1,993.3 | 25 | .0181 |
| 5 | 18°58′ N. | 168°27′ W. | Mar. 11 | 0207–0235 | 1,941.0 | 17 | .0145 |
| 6 | 20°59′ N. | 167°07′ W. | do | 1913–1940 | 1,938.5 | 24 | .0145 |

[1] +10 zone time on stations 1 to 49, +11 zone time on stations 50 to 106.

Cruises 2, 5, and 8 were combined hydrographic and plankton cruises, and thus provide information on currents, temperatures, oxygen, and inorganic phosphate for comparison with the zooplankton abundance. On cruise 7, which was a combination longline-fishing and plankton cruise, subsurface (to approximately 800 feet) and surface temperatures were obtained at all plankton stations.

### SAMPLING IN THE FIELD

Although the primary aim of our plankton sampling was to obtain information on the zooplankton populations of different ocean areas, a secondary objective was to collect tuna eggs and larvae for use in the study of the spawning habits of tuna. Sampling methods and procedures were therefore designed to contribute information toward both objectives.

Plankton nets of the following three types were used on cruise 2:

1. 1-meter (mouth diameter) net with body (front and middle sections) of 30xxx silk grit gauze (width of apertures 0.65 mm.), rear section and bag of 56xxx silk grit gauze (width of apertures 0.31 mm.);
2. 1-meter net with body of 18xxx grit gauze (width of apertures 1.3 mm.), rear section and bag of 30xxx grit gauze; and
3. 2-meter net with body of 3/16 inch square cotton mesh, rear section and bag of 18xxx grit gauze.

The results of this cruise showed that th 1-meter net with a body of 18xxx grit gauze r tained considerably less zooplankton than th 30xxx net; the 2-meter net captured almost n zooplankton and few fish or other large form Some preliminary tows, the results of which a not included in this report, indicated that nets c 56xxx and 72xxx grit gauze retained the larg phytoplankton as well as zooplankton, thus mal ing analysis of the sample more difficult. On th basis of this experimentation and a review of met ods used by other investigators for samplin zooplankton, we adopted the 1-meter, 30xxx n (fig. 2) as being the best suited for our purpose Nets of this type were employed exclusively c cruises 5, 7, and 8.

In this study, sampling was limited to a sing tow at each station. For this reason, we chose use, for the greater part, an oblique tow (surfa to 200 meters to surface) of approximately minutes' duration. On cruise 2, both oblique an surface tows were used, but on all subsequen cruises the oblique tow was the only type em ployed.

The merits of the oblique haul have been we demonstrated by Winsor and Clark (1940). The obtained a percentage standard deviation (coeff cient of variation) for a single observation 31 percent for oblique hauls, 53 percent for vert




DESCRIPTION OF PLANKTON NET SECTIONS

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| MATERIAL | GALV. IRON RING | 12 OZ. CANVAS | 30 XXX GRIT GAUGE | 3/16" BAIT NETTING | 10 OZ. CANVAS | 30 XXX GRIT GAUGE | 10 OZ. CANVAS | 56 XXX GRIT GAUGE | 10 OZ. CANVAS | BUCKET COUPLING | 10 OZ. CANVAS | 56 XXX GRI GAUGE |
| DIAMETER | 39.4" | 39.4" | 39.4" | 40" | 39.4" | 39.4" - 8" | 8" | 8"-4" | 4" | 4" | 4" | 4" |
| LENGTH | 7/8" THICK | 12" | 38" | 48" | 10" | 110" | 3" | 18" | 2 1/2" | 2 1/2" | 2 1/2" | 9" |

FIGURE 2.—Diagram and description of the 1-meter net used in this investigation, showing general construction the net and method of attachment to the weight and towing lines.

l hauls, and 124 percent for horizontal hauls; d they concluded that oblique hauls gave more liable and consistent results than vertical or rizontal hauls. The more erratic nature of the rface haul, as compared with the oblique, is evi- nt from figure 4, showing the results of cruise 2, which both types of haul were used.

That the oblique and surface hauls made on this uise were not greatly different, however, is in- cated by a comparison of the volumes (cc./m.$^3$) tained at 12 pairs of stations (table 1, A and , the members of each pair occurring at approx- ately the same latitude, but at a different longi- de (fig. 1, A). It may be shown that the mean fference (0.00575 cc./m.$^3$) between the paired uls does not differ significantly from 0 ($P$=0.4). herefore, the few surface hauls made on cruise 2 ve been included in this report but have been itted from the statistical analysis dealing with urces of variation and correlations with environ- ental factors.

On cruises 2, 5, and 8, stations were visited nsecutively regardless of the time of day or ght. Because of this practice, the effects of urnal migration must be considered in an evalu- ion of zooplankton abundance at any place and ne. Initially we had hoped, through the use the oblique tow to 200 meters' depth, to nullify a large extent differences in the samples caused vertical movements of the plankton. That we d not succeed is indicated by close perusal of e tables of data, particularly for cruise 5 (table , and cruise 8 (table 4), comparing successive y and night hauls. It will be seen that usually e night hauls produced a higher volume than the y hauls. This tendency will be discussed later the section on sources of variation in zooplank- n volumes.

Another problem that adds to the difficulty of timating zooplankton abundance is the uneven stribution of plankton organisms. This has en referred to by many planktologists and is mphasized by Haeckel (1890), Herdman (1923), ardiner (1931), Hardy (1936), Wilson (1942), iley and Bumpus (1946), Sears (1950), and hers. By our method of straining a large vol- ne of water, averaging over 1,000 cubic meters r haul, and of sampling in uniform fashion from e surface to 200 meters' depth, it is assumed that e variation in catch due to the uneven distribu- on of organisms is minimized.

The amount of water strained during each haul was measured by a flow meter suspended in the mouth of the net. Each flow meter was calibrated by towing it over a measured course at approximately the same speed used in making the plankton hauls. The flow meters were calibrated before and after each cruise, and the average of these calibrations was used to compute the volume of water strained in cubic meters for each haul during that cruise. Within a limited range of towing speeds the number of revolutions registered by the meter indicates the length of the water column passing through the net; multiplying this length by the area of the mouth of the net gives an estimate of the water volume strained.

There has never been evidence of clogging on any of the tows, possibly because of the relatively coarse mesh used in the nets and the general paucity of plankton.

In making the tow, the net and a 75-pound streamlined weight were attached to the cable, which was paid out slowly at uniform speed. As the net was lowered, the length of wire out and the angle of stray were recorded at 2-minute intervals. As soon as a calculated depth of 200 meters was reached, the net was retrieved at a slow, uniform speed. The wire angle and the length of wire out were again recorded at 2-minute intervals. At a towing speed of about 2 knots, an oblique tow to a depth of 200 meters and return required about 30 minutes. A graph of depth reached plotted against time (fig. 3) for 3 tows made on cruise 5, shows that, for practical purposes, equal amounts of time are spent at all depths; i. e., assuming the towing wire represents a straight line in the water, the net strains approximately the same amount of water for each meter of depth passed through. It is recognized that the towing wire does not actually describe a straight line during the tow, but the error caused by a slight curve is small.

When the net reached the surface at the end of a tow, it was lifted out of the water, suspended vertically from a boom, and washed down with a hose. The plankton bucket was then detached, and its contents were washed into an enameled pan. Next, the sample was transferred to a 1-quart fruit jar, and sufficient formalin was added to approximate a 10-percent solution. The formalin was neutralized with borax. A completed label was placed in the jar.




FIGURE 3.—Graphs of three typical tows (stations 1, 12, and 17) of cruise 5, showing depth estimated from wire a
and wire out.

## TREATMENT OF SAMPLES

The zooplankton collections received the following treatment in the laboratory:

1. All fish eggs and larvae were removed. In l cases these have amounted to a negligible fraction of the sample and were omitted from the olume measurement.

2. Several portions of the remaining sample ere examined microscopically, and the various roups of organisms were identified at least to the rder. Where possible without great expenditure f time and effort, identifications were made to the enus and species. A list of constituents was thus mpiled for each sample.

3. All organisms whose longest dimension was reater than 5 centimeters were removed from the mple. Such organisms occurred infrequently nd were not considered in the analysis.

4. By means of a splitting chamber the remaining sample was then divided into halves; one-half as labeled and placed in a reference collection, ne other was used for the organism counts and isplacement-volume determinations.

5. In making the count, a given portion (usually fourth, occasionally a half or an eighth) of the alf reserved for this purpose, was placed in a 5 by 20 cm. counting chamber, thoroughly mixed, nd distributed as evenly as possible over the ntire cell area. Organisms between 2 and 5 cm. their longest dimension were enumerated in ounting-chamber fields 5 cm. square, without agnification. For counting smaller organisms 0.5 millimeter to 2.0 cm. in longest dimension), ne counting cell was placed over a Wolffhuegel late under a binocular microscope. Organisms this size category were counted in fields 1 cm. quare on the Wolffhuegel plate. Organisms less an 0.5 mm. long were identified but not counted, nce the mesh of the nets employed was not sufficiently fine to catch these forms quantitatively. counting organisms of all sizes, either (a) ough fields were counted to yield a minimum ount of 100 for each type of organism, or (b) 10 elds were counted, whichever was reached first. he estimated number of zooplankters of each ecies or category in the total sample (minus fish gs and larvae and organisms larger than 5 cm.) as computed by use of the following formula:

$$E=\frac{C\cdot A}{f\cdot a\cdot n}$$

where

$E$ = estimated number in total sample
$C$ = counted number
$A$ = area of cell
$f$ = fraction of total sample in the counting cell
$a$ = area of field
$n$ = number of fields counted

The estimated total number of zooplankters in the sample equals the sum of $E$ values, i. e., the sum of the estimated numbers for each type or group of organisms. The estimated number per cubic meter of water was obtained by dividing the estimated total number in the sample by the cubic meters of water strained.

6. To measure the displacement volume, the half of the sample used for the organism counts was poured into a draining sock of 56xxx grit gauze to filter off the preserving liquid. The drained plankton was then placed in a 50- or 100-milliliter graduated cylinder, depending upon the size of the sample. By means of a burette a known volume of water was added to the drained plankton. The difference between the volume of the plankton plus the added liquid and the volume of liquid alone is the displacement volume or "wet volume" of the plankton half-sample. This figure was doubled to obtain the computed volume of the entire original sample (minus fish eggs and larvae and organisms larger than 5 cm.). When divided by cubic meters of water strained, this gave the volume (in cubic centimeters) of zooplankton per cubic meter of water.

The plankton counts given in this report, therefore, include those organisms between 0.5 mm. and 5 cm., longest dimensions; the volume measurements include all organisms in the sample less than 5 cm. in length.

Almost all stations occupied on the four cruises were in areas where the depth of water is 2,000 to 3,000 fathoms (roughly 4,000 to 6,000 meters). By sampling to a calculated depth of 200 meters (the actual depth attained was probably within 10 percent of this) we have done little more than "scratch the surface." No information was obtained on plankton abundance and distribution below the sampled depth. Therefore, we believe

that our results are most properly expressed in terms of organisms per cubic meter in the upper 200 meters of water. If an estimate is desired of the areal abundance, i. e., the quantity of zooplankton in a column of water 200 meters in height and 1 meter square in cross section, it may be obtained by multiplying by 200 the numbers and volumes of zooplankton per cubic meter given in tables 1, 2, 3, and 4.

## COMPOSITION OF THE ZOOPLANKTON

Tables 5, 6, 7, and 8 show for each cruise and station the percentage composition by number of the 6 major constituents of the zooplankton. It is evident that copepods—by number—were consistently the most important constituent. The chaetognaths usually were second in rank, followed by the tunicates, euphausiids, siphonophores, and foraminifers. At several stations other groups, such as radiolarians, annelids, amphipods, ostracods, and pteropods occurred in considerable numbers, but on the average these animals formed a very small percentage of the collections. As might be expected, the results vary to some extent from station to station; but when the data are summarized, as in table 9, there is revealed a marked and surprising uniformity in composition for the different longitudes and cruises. The percentages given are computed from the sums of estimated numbers for all stations. Percentages by volume for the separate groups were not determined.

The 210 samples include only a few instances of swarming: Collections taken at stations 7 and 9, cruise 5, contained unusually high percentages (60 and 58 percent) of foraminifers; the collection made at station 2, cruise 7, contained a high and unusual percentage (51 percent) of hyperid amphipods. In each case there was no marked change in number for the other major constituents.

As all of our plankton stations were located in the open ocean with very few within 100 miles of land, the collections consisted primarily of such forms as are permanently planktonic throughout their lives (holoplankton), and contained very few transitory young and larval stages (meroplankton) of bottom-dwelling forms such as echinoderms, crabs, and clams.

TABLE 5.—*Cruise 2: Percentage, by number, of six ma constituents of zooplankton collections*

[Values less than 1 percent omitted]

| Station No. | Copepoda | Chaetognatha | Tunicata | Euphausiacea | Siphonophora | Foraminifera | Mis lane |
|---|---|---|---|---|---|---|---|
| A. Oblique tows to 200 m. (1-meter net, 30xxx mesh): | | | | | | | |
| 1 | 47 | 17 | 13 | 1 | 4 | | |
| 3 | 35 | 19 | 31 | 4 | 2 | 1 | |
| 7 | 64 | 10 | 1 | 6 | 2 | | |
| 9 | 65 | 14 | 4 | 6 | 2 | | |
| 11 | 50 | 16 | 11 | 4 | 3 | 1 | |
| 13 | 61 | 10 | 3 | 4 | | 8 | |
| 15 | 62 | 15 | 3 | 6 | 8 | 5 | |
| 17 | 61 | 10 | 6 | 3 | 4 | 11 | |
| 19 | 55 | 19 | 11 | 4 | 2 | 3 | |
| 21 | 62 | 22 | 3 | 4 | | 1 | |
| 23 | 66 | 16 | 2 | 7 | 2 | | |
| 25 | 68 | 16 | 3 | 2 | 4 | 3 | |
| B. Oblique tows to 200 m. (2-meter net, 3/16-inch mesh, rear section 18xxx mesh): | | | | | | | |
| 2 | 62 | 4 | 14 | 2 | 2 | | |
| 6 | 34 | | 6 | 6 | 22 | | |
| 8 | 56 | 6 | 8 | 6 | 8 | | |
| 10 | 20 | 38 | 20 | | 17 | 1 | |
| 12 | 47 | 16 | 11 | 6 | 10 | | |
| 14 | 44 | 13 | 9 | 4 | 9 | 1 | |
| 16 | 36 | 15 | 8 | 13 | 10 | | |
| 18 | 67 | 12 | 3 | 4 | 5 | | |
| 20 | 62 | 17 | 9 | 4 | 2 | | |
| 22 | 67 | 20 | | 3 | 5 | | |
| 24 | 51 | 19 | | 3 | 19 | | |
| 26 | 19 | 21 | 4 | 12 | 8 | | |
| C. Surface tows (1-meter net, 30xxx mesh): | | | | | | | |
| 30 | 79 | 11 | 3 | 3 | | | |
| 32 | 65 | 12 | 9 | 6 | 3 | | |
| 34 | 53 | 10 | 10 | 14 | 2 | 1 | |
| 36 | 54 | 21 | 6 | 10 | | 6 | |
| 38 | 43 | 15 | 15 | 10 | 1 | 4 | |
| 40 | 51 | 26 | 16 | | 2 | 2 | |
| 42 | 51 | 9 | 24 | 2 | 5 | 3 | |
| 44 | 70 | 5 | 7 | 6 | 3 | 1 | |
| 46 | 76 | 8 | 2 | 5 | 2 | 5 | |
| 48 | 68 | 10 | 2 | 2 | 1 | 9 | |
| 50 | 47 | 19 | 21 | 1 | 3 | 6 | |
| 52 | 50 | 10 | 12 | 2 | 4 | 1 | |
| D. Surface tows (1-meter net, 18xxx mesh): | | | | | | | |
| 31 | 50 | 14 | | 26 | 3 | | |
| 33 | 33 | 43 | 12 | 2 | 8 | | |
| 35 | 24 | 37 | 18 | | 10 | | |
| 37 | 33 | 18 | 9 | 22 | 9 | | |
| 39 | 28 | 29 | 2 | 25 | 9 | | |
| 41 | 57 | 6 | 5 | 25 | 4 | | |
| 43 | 47 | 28 | 8 | | 6 | | |
| 45 | 29 | 5 | 8 | | 12 | | |
| 47 | 47 | 6 | 8 | | 24 | | |
| 49 | 52 | 12 | 2 | 29 | 3 | 2 | |
| 51 | 16 | 25 | 7 | | 13 | 1 | |
| 53 | 37 | 11 | 8 | | 12 | | |

TABLE 6.—*Cruise 5: Percentage, by number, of six ma constituents of zooplankton collections*

[Values less than 1 percent omitted]

| Station No. | Copepoda | Chaetognatha | Tunicata | Euphausiacea | Siphonophora | Foraminifera | Mi lan |
|---|---|---|---|---|---|---|---|
| 1 | 53 | 4 | 9 | 4 | | 10 | |
| 2 | 44 | 4 | 6 | 9 | | 6 | |
| 3 | 58 | 6 | 8 | 5 | 3 | 2 | |
| 4 | 35 | 7 | 11 | 1 | 2 | 21 | |
| 5 | 46 | 5 | 8 | 6 | 3 | 12 | |
| 6 | 41 | 7 | 3 | 12 | 5 | 20 | |
| 7 | 25 | 4 | 4 | | 2 | 60 | |
| 8 | 52 | 1 | 2 | | 3 | 34 | |
| 9 | 22 | 6 | 3 | 2 | 1 | 58 | |
| 10 | 66 | 11 | 6 | 2 | | 9 | |
| 11 | 62 | 12 | 2 | 4 | 4 | 7 | |

TABLE 6.—*Cruise 5: Percentage, by number, of six major constituents of zooplankton collections*—Continued

| Station No. | Copepoda | Chaetognatha | Tunicata | Euphausiacea | Siphonophora | Foraminifera | Miscellaneous |
|---|---|---|---|---|---|---|---|
| 2 | 73 | 8 | 3 | 1 | 8 | ------- | 6 |
| 3 | 61 | 14 | 6 | 4 | 1 | 2 | 12 |
| 4 | 77 | 3 | 3 | 3 | 3 | ------- | 11 |
| 5 | 59 | 9 | 5 | 2 | 7 | 11 | 7 |
| 6 | 52 | 6 | 5 | 2 | 8 | 3 | 24 |
| 7 | 67 | 5 | 3 | 4 | 4 | 2 | 15 |
| 8 | 55 | 11 | 8 | 3 | 10 | 2 | 11 |
| 9 | 68 | 4 | 8 | 8 | 5 | 2 | 5 |
| 0 | 60 | 9 | 8 | 6 | 4 | 2 | 11 |
| 1 | 61 | 11 | 6 | 6 | 1 | ------- | 14 |
| 2 | 56 | 8 | 5 | 6 | 10 | 5 | 10 |
| 3 | 60 | 8 | 5 | 4 | 7 | 7 | 9 |
| 4 | 57 | 10 | 3 | 8 | 4 | 1 | 17 |
| 5 | 54 | 12 | 9 | 5 | 7 | 2 | 11 |
| 6 | 67 | 13 | 7 | 2 | 4 | ------- | 7 |
| 7 | 61 | 16 | 4 | 8 | 2 | ------- | 8 |
| 8 | 70 | 9 | 7 | 4 | 2 | ------- | 8 |
| 9 | 51 | 10 | 5 | 4 | ------- | 9 | 20 |
| 0 | 47 | 10 | 7 | 3 | 2 | 6 | 25 |
| 1 | 56 | 9 | 4 | 7 | 2 | 8 | 14 |
| 2 | 56 | 12 | 8 | 4 | 1 | 3 | 16 |
| 3 | 55 | 14 | 4 | 6 | 3 | 4 | 14 |
| 4 | 53 | 17 | 4 | 2 | 2 | 9 | 13 |
| 5 | 35 | 16 | 3 | 3 | 4 | 26 | 13 |
| 6 | 39 | 9 | 7 | 3 | 5 | 12 | 25 |
| 7 | 50 | 8 | 7 | 4 | 4 | 11 | 16 |
| 8 | 52 | 5 | 6 | 3 | 4 | 13 | 17 |
| 9 | 41 | 5 | 8 | 6 | 4 | 19 | 17 |
| 0 | 56 | 14 | 4 | 2 | 2 | 9 | 13 |
| 1 | 69 | 6 | 1 | 1 | 5 | 10 | 8 |
| 2 | 49 | 11 | 4 | 6 | 2 | 11 | 17 |
| 3 | 58 | 18 | 4 | 2 | 3 | 4 | 11 |
| 4 | 53 | 15 | 6 | 3 | 4 | 8 | 11 |
| 5 | 61 | 13 | 2 | 2 | 3 | 11 | 8 |
| 6 | 63 | 9 | 9 | 6 | 2 | 3 | 8 |
| 7 | 65 | 11 | 4 | 2 | 3 | 11 | 4 |
| 8 | 35 | 6 | 31 | 1 | ------- | 21 | 6 |
| 9 | 58 | 2 | 5 | 1 | 4 | 10 | 20 |
| 0 | 58 | 8 | 6 | 5 | 3 | 2 | 18 |
| 1 | 51 | 4 | 10 | 5 | 5 | 3 | 22 |

TABLE 7.—*Cruise 7: Percentage, by number, of six major constituents of zooplankton collections*

[Values less than 1 percent omitted]

| Station No. | Copepoda | Chaetognatha | Tunicata | Euphausiacea | Siphonophora | Foraminifera | Miscellaneous |
|---|---|---|---|---|---|---|---|
| | 47 | 6 | 4 | 4 | 5 | 7 | 27 |
| | 24 | 5 | 2 | 5 | 2 | 2 | 60 |
| | 51 | 13 | 4 | | 8 | 6 | 14 |
| | 51 | 12 | 16 | 4 | 1 | 6 | 10 |
| | 57 | 7 | 10 | 3 | 5 | 7 | 11 |
| | 59 | 21 | 4 | 2 | ------- | 11 | 3 |
| | 56 | 12 | 3 | 4 | 4 | 6 | 15 |
| | 68 | 11 | 5 | 5 | 4 | 2 | 5 |
| | 68 | 9 | 5 | 2 | 2 | 2 | 12 |
| | 78 | 6 | 4 | ------- | 1 | 2 | 8 |
| | 60 | 8 | 14 | ------- | 3 | 2 | 12 |
| | 75 | 6 | 6 | 3 | 4 | 1 | 5 |
| | 63 | 8 | 6 | 2 | 10 | 3 | 8 |
| | 78 | 4 | 3 | 2 | 2 | 2 | 9 |
| | 52 | 24 | 4 | 4 | 3 | 2 | 11 |
| | 63 | 11 | 2 | 2 | 4 | 5 | 13 |
| | 55 | 17 | 4 | 4 | 4 | 6 | 10 |
| | 61 | 12 | 4 | 2 | 7 | 3 | 11 |
| | 60 | 16 | 2 | 1 | 5 | 2 | 14 |
| | 64 | 14 | 2 | 3 | 7 | ------- | 10 |
| | 60 | 12 | 7 | 4 | 6 | 1 | 10 |
| | 66 | 8 | 2 | 3 | 6 | ------- | 15 |
| | 62 | 14 | 5 | 5 | 3 | 1 | 10 |
| | 62 | 6 | 11 | 5 | 4 | 2 | 10 |

TABLE 8.—*Cruise 8: Percentage, by number, of six major constituents of zooplankton collections*

[Values less than 1 percent omitted]

| Station No. | Copepoda | Chaetognatha | Tunicata | Euphausiacea | Siphonophora | Foraminifera | Miscellaneous |
|---|---|---|---|---|---|---|---|
| 1 | 36 | 10 | 11 | 9 | 6 | ------- | 27 |
| 2 | 52 | 11 | 10 | 3 | 8 | ------- | 16 |
| 4 | 49 | 10 | 12 | 2 | 10 | 13 | 4 |
| 6 | 70 | 6 | 4 | 4 | 1 | 7 | 8 |
| 7 | 58 | 11 | 2 | 3 | 3 | 11 | 12 |
| 8 | 64 | 9 | 3 | 2 | 6 | 8 | 8 |
| 9 | 60 | 4 | 19 | 3 | 3 | 4 | 7 |
| 10 | 66 | 4 | 6 | 3 | 9 | 6 | 6 |
| 11 | 58 | 4 | 4 | 6 | 2 | 12 | 14 |
| 12 | 63 | 5 | 1 | 10 | 4 | 4 | 13 |
| 13 | 59 | 10 | 4 | 7 | 4 | 3 | 13 |
| 14 | 56 | 16 | 2 | 11 | 2 | 2 | 11 |
| 15 | 61 | 14 | 2 | 7 | 1 | 2 | 13 |
| 16 | 58 | 12 | 4 | 4 | 7 | 4 | 11 |
| 17 | 49 | 16 | 3 | 5 | 6 | 7 | 14 |
| 18 | 65 | 10 | 6 | 6 | 3 | ------- | 10 |
| 19 | 58 | 11 | 4 | 8 | 3 | 2 | 14 |
| 20 | 54 | 14 | 4 | 3 | 6 | 9 | 10 |
| 21 | 52 | 16 | 2 | 13 | 4 | 3 | 10 |
| 22 | 59 | 20 | 9 | 3 | ------- | 3 | 6 |
| 23 | 63 | 21 | 3 | 3 | ------- | 3 | 6 |
| 24 | 51 | 17 | 2 | 11 | 3 | 6 | 10 |
| 25 | 59 | 17 | 12 | 4 | 2 | 4 | 2 |
| 26 | 66 | 20 | 8 | ------- | 3 | ------- | 3 |
| 32 | 64 | 8 | 4 | 8 | 4 | ------- | 12 |
| 33 | 56 | 22 | 8 | 3 | 4 | ------- | 7 |
| 34 | 64 | 14 | 4 | 6 | 5 | ------- | 7 |
| 35 | 55 | 16 | 6 | 7 | 6 | ------- | 9 |
| 36 | 57 | 19 | 4 | 3 | 4 | 2 | 11 |
| 37 | 50 | 22 | 2 | 10 | 5 | 1 | 10 |
| 42 | 68 | 8 | ------- | 11 | 3 | ------- | 10 |
| 43 | 74 | 11 | 2 | 4 | 1 | ------- | 8 |
| 44 | 66 | 12 | 3 | 8 | 1 | ------- | 10 |
| 45 | 56 | 19 | 4 | 9 | 2 | ------- | 10 |
| 46 | 59 | 22 | ------- | 10 | 4 | ------- | 4 |
| 47 | 59 | 10 | 10 | 5 | 2 | ------- | 13 |
| 52 | 62 | 17 | 7 | 1 | 4 | 3 | 6 |
| 53 | 62 | 12 | 5 | 6 | 3 | 2 | 10 |
| 54 | 55 | 17 | 6 | 6 | 2 | ------- | 13 |
| 55 | 54 | 22 | 7 | 2 | 1 | 4 | 10 |
| 56 | 51 | 13 | 8 | 2 | 3 | ------- | 22 |
| 57 | 56 | 20 | 2 | 11 | 2 | 1 | 8 |
| 62 | 64 | 13 | 3 | 2 | 4 | 2 | 12 |
| 63 | 58 | 14 | 5 | 9 | 3 | 3 | 8 |
| 64 | 59 | 12 | 3 | 9 | 4 | 4 | 9 |
| 65 | 63 | 21 | 6 | 2 | 1 | ------- | 6 |
| 66 | 61 | 6 | 7 | 7 | 5 | 4 | 10 |
| 67 | 56 | 19 | 5 | 2 | 2 | 2 | 14 |
| 68 | 47 | 18 | 2 | 2 | 4 | 7 | 20 |
| 69 | 67 | 8 | 3 | 6 | 3 | 3 | 10 |
| 70 | 62 | 18 | 4 | 1 | 4 | 2 | 9 |
| 71 | 53 | 10 | 4 | 7 | 8 | 3 | 15 |
| 72 | 55 | 11 | 2 | 6 | 9 | 2 | 15 |
| 73 | 60 | 22 | 8 | 2 | 1 | 2 | 5 |
| 74 | 55 | 24 | 10 | 6 | 2 | ------- | 3 |
| 75 | 67 | 11 | 1 | 8 | 2 | ------- | 10 |
| 76 | 51 | 24 | 9 | 5 | 3 | ------- | 8 |
| 77 | 43 | 6 | 5 | 9 | 14 | 1 | 22 |
| 78 | 55 | 7 | 6 | 1 | 11 | 4 | 16 |
| 79 | 66 | 12 | 5 | 3 | 7 | 1 | 6 |
| 80 | 74 | 7 | 3 | 2 | 3 | ------- | 10 |
| 81 | 65 | 16 | 5 | 5 | 3 | ------- | 6 |
| 82 | 74 | 10 | 5 | 3 | 2 | ------- | 6 |
| 83 | 59 | 19 | ------- | 9 | 2 | ------- | 11 |
| 84 | 68 | 10 | 6 | 1 | 2 | 2 | 11 |
| 85 | 51 | 19 | 12 | 2 | 10 | ------- | 5 |
| 86 | 55 | 27 | 4 | 6 | ------- | ------- | 8 |
| 87 | 54 | 19 | 5 | 6 | 5 | 1 | 10 |
| 88 | 51 | 17 | 5 | 4 | 6 | 1 | 16 |
| 89 | 49 | 17 | 7 | 3 | 4 | 5 | 15 |
| 90 | 60 | 10 | 5 | 6 | 3 | 3 | 13 |
| 91 | 46 | 15 | 10 | 3 | 5 | 2 | 19 |
| 92 | 47 | 17 | 4 | 7 | 9 | 2 | 14 |
| 93 | 70 | 10 | 1 | 5 | 4 | ------- | 9 |
| 94 | 56 | 18 | 7 | 4 | 5 | 4 | 6 |
| 95 | 72 | 6 | 3 | 4 | 3 | 2 | 10 |
| 96 | 72 | 7 | 7 | 7 | ------- | ------- | 5 |
| 97 | 53 | 17 | 10 | 2 | 4 | 8 | 6 |
| 98 | 59 | 9 | 10 | 5 | 4 | ------- | 12 |
| 99 | 55 | 13 | 6 | 5 | 4 | 4 | 13 |
| 100 | 43 | 8 | 16 | 1 | 9 | 2 | 21 |
| 101 | 57 | 9 | 11 | 4 | 8 | ------- | 10 |
| 102 | 54 | 14 | 1 | 9 | 2 | ------- | 19 |
| 103 | 67 | 2 | 9 | 3 | 6 | ------- | 13 |
| 104 | 67 | 11 | 6 | 1 | 5 | ------- | 10 |
| 105 | 70 | 4 | 4 | 3 | 8 | ------- | 11 |
| 106 | 49 | 7 | 15 | 9 | 5 | ------- | 15 |

TABLE 9.—*Percentage composition, by number, of the six major constituents of the zooplankton collections, by cruise an longitude*

| Major constituents | Cruise 2 Jan.–Feb. 1950 | | | Cruise 5 June–Aug. 1950 | | | Cruise 7 Oct.–Nov. 1950 | Cruise 8 Jan.–Mar. 1951 | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 172° W.[1] | 158° W.[2] | Total | 172° W.[3] | 158° W. | Total | 158° W. | 172° W.[4] | 158° W.[5] | Total[6] |
| Copepoda | 58.9 | 56.2 | 57.4 | 56.3 | 53.0 | 54.9 | 60.2 | 58.6 | 58.0 | 58. |
| Chaetognatha | 16.7 | 14.0 | 15.2 | 8.3 | 10.0 | 9.0 | 12.0 | 12.4 | 12.2 | 13. |
| Tunicata | 5.6 | 8.2 | 7.1 | 5.6 | 5.6 | 5.6 | 4.7 | 6.1 | 5.3 | 5. |
| Euphausiacea | 3.9 | 9.1 | 6.8 | 4.9 | 3.5 | 4.3 | 3.0 | 4.6 | 5.9 | 5. |
| Siphonophora | 4.2 | 2.9 | 3.5 | 5.0 | 3.4 | 4.3 | 4.3 | 5.2 | 4.0 | 4. |
| Foraminifera | 2.5 | 1.3 | 1.8 | 7.7 | 9.9 | 8.6 | 3.0 | 1.6 | 4.8 | 2. |
| Miscellaneous | 8.2 | 8.3 | 8.2 | 12.2 | 14.6 | 13.3 | 12.8 | 11.5 | 9.8 | 10. |

[1] Includes stations 1 to 26.
[2] Includes stations 30 to 53.
[3] Includes stations 1 to 27.
[4] Includes stations 77 to 106.
[5] Includes stations 1 to 26.
[6] Includes stations 1 to 106, all longitudes.

The uniform composition of the zooplankton, as shown by our measurements, indicates the existence of a stable ecological balance among the various components of the plankton and between them and their environment. This uniformity, or stability, is doubtless the result of a complex interaction of factors. The physical and chemical factors of the environment throughout the range sampled are rather homogeneous and, perhaps while capable of effecting differences in total abundance of organisms, do not provide sufficient variations to promote major differences in the composition of populations. The relative absence of irregularly occurring swarms of larval and adult forms may account for some of the uniformity. It is reasonable to assume that, despite the great species differences that may occur (Wilson 1942), there is a particular niche in this ecosystem of the tropical and subtropical Pacific which will support a certain number of copepods, and another which will accommodate a certain number of chaetognaths, and so forth. The uniformity of the data on plankton composition also argues that the collecting method and its uniformity of application were appropriate for producing repeatable results.

## ZOOPLANKTON AS FOOD

It is generally recognized that neither an enumeration of organisms present nor a total-volume measurement shows the actual food value of a plankton sample. Food value could be estimated by chemical analysis of each sample, but this procedure is hardly practical when large numbers of samples must be examined. Sufficient work has been done on the chemical composition of the major zooplankton types to show that they vary widely in nutritive value among types and eve within types for different localities.

According to Bigelow and Sears (1939), th separation of the crustaceans and chaetognath from the other types of zooplankton permits a approximate division of the zooplankton into (1 more nutritive forms which may be important fish food and (2) forms of little or no nutriti value, such as the tunicates and siphonophore In our collections, the crustacean-chaetognat group averages 70 to 80 percent by number.

Nakai (1942) has shown that plankton anima from the southern part of the Sea of Japan ge erally contain less fat than those of northern area The inference is made that in the warmer wate to the south, the scarcity of phytoplankton, pa ticularly of diatoms with their rich oil reserve prevents the accumulation of fat by the zooplan ton. Clarke (1940) found that in the weste Atlantic the plankton of coastal water had higher percentage of organic matter than that continental-slope or Sargasso-Sea water. Su tropical plankton in general had a low orga content. It is possible, therefore, that wh three-fourths by number, and a smaller fracti by volume, of the zooplankton of the central P cific is theoretically nutritious, its actual fo value in calories may be less than for similar o ganisms of higher latitudes.

The Pacific Oceanic Fishery Investigations conducting a study of the food of tunas. T results to date show that a variety of zooplankt forms are utilized directly as food by these fi For example, representatives of the followi groups have been captured in our plankton n and have also been found among the stomach c

ents of maturing and adult yellowfin tuna (Reintjes and King 1953):

| | |
|---|---|
| ysids | brachyuran larvae |
| uphausiids | heteropods |
| mphipods | pteropods |
| tomatopod larvae | cephalopod young |
| hrimps | tunicates |
| alinurid larvae | fish young |

References have been previously cited (p. 111)

emphasizing the importance of plankton in the food of juvenile tunas.

## ABUNDANCE OF THE ZOOPLANKTON

Estimated numbers and volumes of zooplankton, times of sampling, and amounts of water strained, are given in tables 1, 2, 3, and 4. Variations in abundance with latitude are demonstrated by the histograms of figures 4, 5, 6, 7, and 8. The results are briefly summarized in tables 10 and 11.

TABLE 10.—*Areas of greatest abundance of zooplankton, by cruise and longitude*

| Zooplankton | Cruise 2 | | Cruise 5 | | Cruise 7 | Cruise 8 | |
|---|---|---|---|---|---|---|---|
| | 172° W. | 158° W. | 172° W. | 158° W. | 158° W. | 172° W. | 158° W. |
| reatest estimated number/m.³ | 109 | 83 | 107 | 93 | 75 | 73 | 90 |
| atitude of greatest estimated number | 0° | 5° S. | 1° S. | 3° N. | 2° N. | 1° N. | 1° S. |
| reatest displacement volume, cc./m.³ | .0826 | .0911 | .0742 | .0880 | .0741 | .0548 | .0666 |
| atitude of greatest displacement volume | 0° | 1° S. | 1° S. | 0° | 2° N.[1] | 2° N. | 5° S. |
| oundaries of rich area (estimated numbers) | 2° N.–0° | 3° N.–5° S. | 4° N.–3° S. | 5° N.–0° | 6° N.–2° N. | 6° N.–1° S. | 1° S.–6° S. |
| oundaries of rich area (displacement volumes) | 2° N.–2° S. | 3° N.–1° S. | 1° N.–3° S. | 5° N.–0° | 6° N.–2° N. | 5° N.–1° S. | 1° S.–5° S. |
| atitudinal range of collection | 24° N.–5° S. | 19° N.–5° S. | 27° N.–5° S. | 21° N.–5° S. | 20½° N.–2° S. | 21°N .–14½° S. | 21° N. –7° S. |

[1] Sample at 19° N. latitude was greatest in volume but was atypical, as it contained a highly unusual number and volume of amphipods. (See table 3.)

TABLE 11.—*Numbers and volumes of zooplankton organisms as related to equatorial currents and latitudes*

Data from the six major sections of cruises 2, 5, and 8, which crossed the Equator on 158° and 172° W. longitude]

| Current | Latitude | Number of observations | Zooplankton organisms | |
|---|---|---|---|---|
| | | | Average number per m.³ | Average volume, cc. per m.³ |
| North Equatorial Current | 27°–18° N. | 14 | 28 | 0.0184 |
| | 17°–11° N. | 27 | 26 | .0176 |
| Countercurrent | 10°–6° N. | 25 | 32 | .0255 |
| South Equatorial Current | 5°–1° N. | 25 | 53 | .0430 |
| | Equator | 5 | 73 | .0605 |
| | 1°–5°S. | 25 | 46 | .0373 |
| | 6°–14° S. | 8 | 34 | .0175 |

In general, at the time and through the range of latitudes sampled (27° N. to 14° S.), the greatest abundance of zooplankton was found in the region of the Equator, between 6° N. and 5° S. latitude. The latitude of peak abundance varied with longitude and cruise. Although not evaluated statistically, there appears in all sections a strikingly parallel variation between the estimated number for each sample and the displacement volume. Because of the disparity in size among the different kinds of zooplankton organisms, numbers are much less meaningful indicators of productivity, or available food, than sample volumes; therefore, our statistical studies have been based solely on volume determinations.

When the varying abundance of zooplankton is reviewed in respect to the presence or absence of a convergence north of the Equator, certain interesting relationships are indicated:

1. When a well-marked convergence is present, the rich zone of zooplankton appears to lie between the Equator and the convergence to the northward (fig. 4, left panel; fig. 5, right panel). The eastern section, cruise 2 (fig. 4, right panel) on which only surface tows were employed, does not conform to this generality. Much of the irregularity in this section, however, is due to differences between samples taken by day and by night.

2. The northern boundary of the rich zone is practically demarcated by the position of the convergence—when it occurs.

3. When a well-marked convergence is lacking, the peaks of abundance—both in number and volume—occur south of the Equator (fig. 5, left panel; fig. 7, right panel).

4. The cruise-8 hydrographic data are not as yet completely processed and the presence or absence of convergences has not been determined for all sections; however, there is no evidence of a convergence along 158° W. longitude (fig. 7, right panel).




FIGURE 4.—Histograms of zooplankton abundance, as estimated numbers and volumes per m.³ of water strained, for cruise-2 stations.




FIGURE 5.—Histograms of zooplankton abundance, as estimated numbers and volumes per $m.^3$ of water strained, for cruise-5 stations.




FIGURE 6.—Histogram of zooplankton abundance, as estimated numbers and volumes per m.³ of water strained, cruise-7 stations.




FIGURE 7.—Histograms of zooplankton abundance, as estimated numbers and volumes per m.³ of water strained, for cruise-8 stations on longitudes 158° and 172° W.




FIGURE 8.—Histograms of zooplankton abundance, as estimated numbers and volumes per m.³ of water strained, for cruise-8 stations located on the short

## EFFECTS OF THE EQUATORIAL CURRENT SYSTEM

The general pattern of the Pacific Equatorial Current system has been described by Sverdrup, Johnson, and Fleming (1942, pp. 708–712). Cromwell (1951 and unpublished ms.[3]), on the basis of recent investigations by this laboratory, suggests certain modifications in the generally accepted ideas pertaining to the details of the circulation. The existence of the equatorial divergence with resultant upwelling of nutrient-rich water, as demonstrated by the *Carnegie* section of 1929 (Sverdrup, et al. 1942), is substantiated by the researches of this laboratory. The presence of a second major divergence at the northern boundary of the Countercurrent has not been confirmed, however. Cromwell concludes, on the basis of the information obtained on three cruises (2, 5, and 8) of the *Hugh M. Smith*, that there is no evidence that any enrichment of the surface layer—as a result of upwelling accompanying a divergence—is occurring at the boundary between the North Equatorial Current and the Countercurrent.

On cruise 2, a sharp temperature discontinuity, together with a local deepening of the thermocline, was found to occur at about 2° N. latitude on the western section and at about 4½° N. latitude in the eastern section (fig. 4). These conditions indicate the presence of a convergence or "front" (Cromwell ms.) in the surface currents. The northern boundary of the South Equatorial Current was discernible at 5½° N. latitude on the western section, but was not well defined on the eastern section. The convergence near 2° N. latitude was entirely within the South Equatorial Current and, therefore, was not associated with the boundary of this current and the Countercurrent as indicated by Sverdrup, et al. (1942, pp. 710, 711) and Arrhenius (1950). A well-defined convergence, as demonstrated by a marked temperature discontinuity and a deepening thermocline, occurred at 4½° N. latitude on the eastern section of cruise 5 (fig. 5) and again was entirely within the South Equatorial Current. An examination of the data from the oceanographic cruises (tables 6 to 8) indicates that the convergence was more pronounced in certain sections than in others and in some it was not strong enough to exhibit the usual signs and may have been entirely lacking, thus permitting the inference that it is shifting and transitory in nature.[4]

Newly upwelled water is at first poorly populated as regards both phytoplankton and zooplankton (Steemann Nielsen 1937). The latter does not benefit directly from the nutrient-rich water but must await the development of a phytoplankton population. In a region of fairly rapid currents, the phytoplankton maximum and the more slowly developing zooplankton maximum would be expected to occur at a considerable distance from each other both in time and space. Steemann Nielsen (1937) reports that in Iceland waters, when conditions for phytoplankton change from bad to good, the capacity for rapid reproduction in phytoplankton enables a few specimens to establish a rich population in 10 days; large quantities of zooplankton do not show up until about a month later. We are not aware of similar population-growth data for the tropics, but we can safely assume that the rate of development here would be considerably more rapid. Nevertheless, it is surprising that in our sampling we found the zooplankton maximum frequently occurring on or very near the site of the upwelling.

Since the days of Hensen and the German Plankton Expedition of 1889, tropical seas have been considered less productive than those of higher latitudes. Explanations for this are usually based on the theory that in the tropics surface heating results in a stable stratification of the sea, thus preventing any vertical mixing by convection which would bring nutrients to the euphotic zone (Delsman 1939; Graham 1941; Sverdrup, et al. 1942, p. 942; Arrhenius 1950). Such thermal stratification does apparently exist throughout much of the tropical, subtropical, and temperate Pacific in areas removed from land influence.

It was learned from the last cruise of the *Carnegie* in 1929, that these generally stable dynamic conditions are disrupted at the Equator by a strong divergence which is accompanied by upwelling. The latter was considered responsible for the enrichment of the surface layers and the production of much higher concentrations of plankton than

[3] Circulation in a meridional plane in the central equatorial Pacific.

[4] The Equatorial Current system is considerably more complex than indicated here. Details of the circulation at the divergence and the convergence and the causal forces involved, are presently being studied by members of this organization and the Scripps Institution of Oceanography.

were found in subtropical and temperate latitudes (Graham 1941). The frequently appearing convergence at 2° to 5° N. latitude also contributes to this apparent greater abundance of life in the equatorial region. Floating organisms with high buoyancy undoubtedly congregate here just as driftwood does along a current rip or other discontinuity zone; zooplankton capable of resisting the downward currents would also tend to concentrate in the surface layers. When a strong convergence has persisted for a sufficient length of time, an area of relatively high-plankton abundance should result, providing rich pasturage for plankton-feeding animals.

The unusual abundance of marine life in the equatorial region has been observed by many expeditions and world travelers while crossing the Pacific. Agassiz and Mayer (1902), in their report on observations and collections of the *Albatross* in the tropical Pacific, state,

> In fact it is evident that pelagic animals are not abundant in regions far from large land masses or where there are no well-defined oceanic currents or counter currents. As soon as one approaches the region of great currents or counter currents, or the coasts of continents and larger islands, the number of animals increases with remarkable suddenness.

Beebe (1926) describes masses of floating debris and the associated wealth of life in a current rip some 200 miles southeast (between 3° and 4° N. latitude) of Cocos Island in the eastern Pacific. Brooks (1934), in reporting his observations on certain relationships between ocean currents and birds, asks why it is that on ocean voyages day after day goes by and no birds or marine life are seen and then suddenly one passes into an area teeming with life. He observed that, as his vessel neared the Equator from the north, bird life increased greatly, as did other marine life such as flying fish, sharks, and whales; then as he crossed the Equator, the zone of abundant life was left behind.

Revelle (1944) points out the agreement between the southern boundary of the Countercurrent and the northern border of the Pacific *Globigerina* ooze area. In reporting on results of the recent Swedish Deep-Sea Expedition, Arrhenius (1950) states in respect to the Pacific,

> The biogenous component and simultaneously the rate of sedimentation increases strongly below the convergence where a biolith, rich in fossils of foraminifera, radiolaria and diatoms is thus deposited. North of th convergence of the Equatorial Counter Current, the sha of the biogenous component in the sedimentation d creases and the fossil-rich biolith turns into a clay, po in fossils and with a low intensity of sedimentation.[5]

As summarized by Herdman (1923),

> It is probable, on the whole, that the distribution an variation of ocean currents have more than latitude temperature alone to do with any observed scantiness tropical plankton. These mighty rivers of the ocean places teem with animal and plant life, and may swee abundance of food from one region to another in th open sea.

## VARIATION IN ZOOPLANKTON ABUNDANCE

The sampling method used in the present stud meets the criteria for randomness in most respect allowing the use of standard methods of statistic analysis. Although the stations were located systematic order, a random sample was probab obtained of the zooplankton population retaine by a meter net with 0.65-mm. apertures from th upper 200 meters of water. There is some doub however, as to the type of distribution of th plankton population. Although our results inc cate a rather uniform, but very dilute distributio of zooplankton, which would perhaps conform a Poisson distribution, the majority of worke have reported clumping and lack of uniformit Snedecor (1946, pp. 42 and 252) states that the r quirement of randomness must be adhered to, b normal distribution of the population is a speci cation that can be considerably relaxed. Of cour there are devices appropriate for analyzing sa ples from nonnormal distributions. One co monly used method involving a transformation the data to logarithms, was employed in certa initial tests of the volume data. The results a conclusions obtained were the same, however, those reached through an analysis of the untra formed data. We assumed, therefore, that t degree of anormality in the zooplankton popu tion would affect our inferences but little, and this report have chosen to examine and base o conclusions upon tests of the untransformed da We wish to emphasize the point that all statis

[5] The convergence here referred to by Arrhenius is that wh theoretically occurs at the southern boundary of the Coun current and, as previously stated, has not been observed by th Investigations. We have found the convergence 2° to 3° so of the current boundary and entirely within the South Equato Current.

al analyses were made on zooplankton volumes ather than on numbers.

An examination of the histograms in figures 4, , 6, 7, and 8 reveals distinct variations in zoo- lankton abundance with latitude; differences be- ween longitudes and between cruises are not so bvious, however. The hydrographic data ob- ained coincident with the zooplankton collections lso show latitudinal variations in inorganic hosphate, surface and subsurface temperature, xygen, and thermocline depth (fig. 9). The narked differences between day and night hauls ave already been referred to in the discussion of ethods. By employing an analysis of variance, e have attempted to determine the significance nd magnitude of these space-time variations in ooplankton abundance; by using correlation and ultiple-regression analyses we have been able to easure the degree of covariation between zoo- lankton and these different environmental actors.

### Effect of time of sampling

Our sampling method was not specifically de- igned to evaluate variations among the zooplank- on volumes resulting from differences between ay and night hauls. An accurate measure of hese differences would have required both day nd night hauls (preferably within a 24-hour pe- iod) at every station. These day-night varia- ions could perhaps have been eliminated by isiting every station at the same time of day or ight. In view of the number of stations and he large area covered, neither procedure was ractical.

Although on cruises 2, 5, and 8 there was, theo- etically, opportunity for the occurrence of equal umbers of day and night collections, it so hap- ened that the day collections outnumbered the ight collections. Rarely, however, were more an 2 day stations or 2 night stations occupied onsecutively.

An estimate of the importance of the day-night ariation, as compared with the latitudinal vari- tion, was obtained from the cruise-5 and cruise-8 ata, using an analysis of variance with two cri- ria of classification (following Snedecor 1946, . 256). We paired, impartially, successive night nd day hauls, where both occurred on the same ngitude and on about the same latitude, omit- ting those stations where two or more day hauls or night hauls were made in succession, and also omitting those stations worked during twilight periods.

An analysis of the cruise-5 volumes (table 12) indicates that for this cruise, there were significant ($P<0.05$) differences between the day and night hauls and also among latitudes. For the cruise-8 volumes (table 13), there were also significant ($P<0.01$) differences between the day and night hauls and among latitudes.

If the means for latitudes are compared using the "least significant difference" calculated accord- ing to the method of Johnson (1950, p. 123; $LSD=(t_{.05})\sqrt{2s^2/k}$), it is evident that for both cruises the means obtained near the Equator are significantly greater than those to the north or south of the Equator. For example, on cruise 8, the mean volume (0.0513 cc./m.$^3$) for 1° and 2° S. latitude differs from the mean volume (0.0178 cc./m.$^3$) for 8° and 9° N. latitude by a difference of 0.0335, which is considerably greater than the least significant difference, 0.0165.

TABLE 12.—*Cruise 5: Analysis of variance of volumes of paired samples taken by day and by night in about the same latitude*

[Two criteria of classification]

| Station Nos. | Latitudes | Volume (cc./m.$^3$) | | Mean |
|---|---|---|---|---|
| | | Day | Night | |
| 1 and 2 | 27° and 25° N | 0.0118 | 0.0271 | 0.0194 |
| 5 and 4 | 19° and 21° N | .0174 | .0254 | .0214 |
| 7 and 8 | 15° and 14° N | .0216 | .0234 | .0225 |
| 12 and 11 | 10° and 11° N | .0252 | .0374 | .0313 |
| 15 and 14 | 7° and 8° N | .0204 | .0203 | .0204 |
| 17 and 16 | 5° and 6° N | .0226 | .0338 | .0282 |
| 18 and 19 | 4° and 3° N | .0385 | .0410 | .0398 |
| 21 and 22 | 1° N. and 0° | .0556 | .0604 | .0580 |
| 24 and 25 | 2° and 3° S | .0452 | .0664 | .0558 |
| 26 and 27 | 4° and 5° S | .0230 | .0315 | .0272 |
| 29 and 30 | 4° and 3° S | .0239 | .0490 | .0370 |
| 32 and 33 | 1° S. and 0° | .0239 | .0880 | .0560 |
| 35 and 36 | 2° and 3° N | .0658 | .0701 | .0680 |
| 38 and 39 | 5° and 6° N | .0691 | .0449 | .0570 |
| 41 and 42 | 8° and 9° N | .0230 | .0531 | .0380 |
| 44 and 45 | 11° and 12° N | .0242 | .0327 | .0284 |
| 47 and 48 | 14° and 15° N | .0201 | .0246 | .0224 |
| 50 and 51 | 19° and 21° N | .0357 | .0289 | .0323 |
| Mean | | .0315 | .0421 | |

Least significant difference for latitudinal means=.0268

| Source of variation | Degrees of freedom | Sum of squares | Mean square | *F* | *P* |
|---|---|---|---|---|---|
| Times | 1 | 0.00101336 | 0.00101336* | 6.28 | <0.05 |
| Latitudes | 17 | .00814675 | .00047922* | 2.97 | <0.05 |
| Discrepancy | 17 | .00274379 | .00016140 | | |
| Total | 35 | .01190390 | | | |

*Indicates a significant ($P<0.05$) mean square value.




FIGURE 9.—Variations in temperature, salinity, oxygen, inorganic phosphate, thermocline depth, and zooplankt volume along 158° and 172° W. long. as found on cruise 5, *Hugh M. Smith*, June–August 1950.

ABLE 13.—*Cruise 8: Analysis of variance of volumes of paired samples taken by day and by night in about the same latitude*

[Two criteria of classification]

| Station Nos. | Latitudes | Volume (cc./m.³) | | Mean |
|---|---|---|---|---|
| | | Day | Night | |
| nd 1 | 19° and 21° N. | 0.0105 | 0.0154 | 0.0130 |
| nd 6 | 14½° and 13° N. | .0236 | .0310 | .0273 |
| nd 9 | 11° and 10° N. | .0093 | .0251 | .0172 |
| and 11 | 9° and 8° N. | .0166 | .0190 | .0178 |
| and 14 | 6° and 5° N. | .0186 | .0174 | .0180 |
| and 16 | 4° and 3° N. | .0118 | .0255 | .0186 |
| and 19 | 1° N. and 0° | .0220 | .0370 | .0295 |
| and 21 | 1° and 2° S. | .0478 | .0558 | .0513 |
| and 24 | 4° and 5° S. | .0178 | .0666 | .0422 |
| and 32 | 6° and 7° N. | .0325 | .0251 | .0288 |
| and 35 | 3° and 4° N. | .0348 | .0340 | .0344 |
| and 44 | 6° and 5° N. | .0126 | .0204 | .0165 |
| and 46 | 4° and 3° N. | .0177 | .0254 | .0216 |
| and 53 | 7° and 6° N. | .0224 | .0313 | .0208 |
| and 54 | 4° and 5° N. | .0118 | .0276 | .0197 |
| and 63 | 7° and 6° N. | .0170 | .0368 | .0269 |
| and 66 | 4° and 3° N. | .0118 | .0359 | .0238 |
| and 69 | 1° S. and 0° | .0172 | .0473 | .0322 |
| and 72 | 4° and 3° S. | .0175 | .0295 | .0235 |
| and 75 | 7° and 6° S. | .0100 | .0332 | .0216 |
| and 79 | 12° and 10° S. | .0115 | .0226 | .0170 |
| and 81 | 6° and 7° S. | .0100 | .0291 | .0196 |
| and 83 | 4° and 5° S. | .0101 | .0202 | .0152 |
| and 87 | 2° and 1° S. | .0109 | .0394 | .0252 |
| and 90 | 1° and 2° N. | .0398 | .0548 | .0473 |
| and 93 | 6° and 5° N. | .0242 | .0369 | .0306 |
| and 98 | 9° and 10° N. | .0094 | .0127 | .0110 |
| and 101 | 12° and 13° N. | .0097 | .0192 | .0144 |
| and 105 | 17° and 19° N. | .0181 | .0145 | .0163 |
| Mean | | .0182 | .0306 | |

Least significant difference for latitudinal means = .0165

| ource of variation | Degrees of freedom | Sum of squares | Mean square | *F* | *P* |
|---|---|---|---|---|---|
| nes | 1 | 0.00225564 | 0.00225564** | 34.60 | <0.01 |
| itudes | 28 | .00553218 | .00019758** | 3.03 | <0.01 |
| crepancy | 28 | .00182531 | .00006519 | | |
| Total | 57 | .00961313 | | | |

*Indicates a highly significant ($P<0.01$) mean square value.

Thus, despite the important variation resulting om diurnal migration of the zooplankton, there ill remain in our data significant differences nong latitudes. Zooplankton populations oc- rring near the Equator are significantly greater abundance than those of adjoining areas.

### riations between cruises, latitudes, and longitudes

To examine differences between cruises and be- een longitudes and to inspect further the differ- ces among latitudes, an analysis of variance with ltiple classification as outlined by Snedecor 946, pp. 304–309) was utilized, employing as ny of the cruise-5 and cruise-8 data as were ailable for similar latitudes and longitudes able 14). The analysis is of the same general pe as that used by Winsor and Clarke (1940) their study of variation in the catch of plank- n nets.

TABLE 14.—*Analysis of variance of volumes (cc./m.³) of zooplankton samples collected on cruises 5 and 8*

[Three criteria of classification]

| Latitude | Series A (172° W. longitude) | | Series B (158° W. longitude) | | Mean |
|---|---|---|---|---|---|
| | Cruise 5 | Cruise 8 | Cruise 5 | Cruise 8 | |
| 21° N. | 0.0254 | 0.0145 | 0.0289 | 0.0154 | 0.0210 |
| 19° N. | .0174 | .0145 | .0357 | .0105 | .0195 |
| 15° N. | .0216 | .0100 | .0246 | .0236 | .0200 |
| 13° N. | .0125 | .0192 | .0158 | .0310 | .0196 |
| 12° N. | .0084 | .0097 | .0327 | .0094 | .0150 |
| 11° N. | .0374 | .0115 | .0242 | .0093 | .0206 |
| 10° N. | .0252 | .0127 | .0225 | .0251 | .0214 |
| 9° N. | .0301 | .0094 | .0531 | .0166 | .0273 |
| 8° N. | .0203 | .0139 | .0230 | .0190 | .0190 |
| 7° N. | .0204 | .0244 | .0258 | .0477 | .0296 |
| 6° N. | .0338 | .0242 | .0449 | .0186 | .0304 |
| 5° N. | .0226 | .0369 | .0691 | .0174 | .0365 |
| 4° N. | .0385 | .0367 | .0783 | .0118 | .0413 |
| 3° N. | .0410 | .0435 | .0701 | .0255 | .0450 |
| 2° N. | .0330 | .0548 | .0658 | .0146 | .0420 |
| 1° N. | .0556 | .0398 | .0347 | .0220 | .0380 |
| 0° | .0604 | .0345 | .0880 | .0370 | .0550 |
| 1° S. | .0742 | .0394 | .0239 | .0478 | .0463 |
| 2° S. | .0452 | .0109 | .0253 | .0558 | .0343 |
| 3° S. | .0664 | .0146 | .0490 | .0257 | .0389 |
| 4° S. | .0230 | .0101 | .0239 | .0178 | .0187 |
| 5° S. | .0315 | .0202 | .0087 | .0666 | .0318 |
| Mean | .0338 | .0230 | .0394 | .0258 | |

Least significant difference for latitudinal means = .0218

| Source of variation | Degrees of freedom | Sum of squares | Mean square | *F* | *P* |
|---|---|---|---|---|---|
| Main effects: | | | | | |
| Series (S) | 1 | 0.00039964 | 0.00039964 | 1.67 | >0.05 |
| Cruises (C) | 1 | .00329280 | .00329280** | 13.79 | <0.01 |
| Latitudes (L) | 21 | .01052548 | .00050121* | 2.10 | <0.02 |
| First-order interactions: | | | | | |
| S×C | 1 | | | | |
| S×L | 21 | | | | |
| C×L | 21 | .01528396 | .00023881 | | |
| Second-order interaction: | | | | | |
| S×C×L | 21 | | | | |
| Total | 87 | .02950188 | | | |

* Indicates a significant ($P<0.05$) mean square value.
** Indicates a highly significant ($P<0.01$) mean square value.

The analysis was first carried out in full to determine the significance of the first-order interactions. As these proved to be nonsignificant, they were pooled (following Kendall 1948, p. 201) with the second-order interaction to give a new sum of squares with 64 degrees of freedom and a new mean square which was then used as the error term for testing the main effects. The following observations may be made from the tests of significance:

1. No significant differences ($P>0.05$) are demonstrated between the means for series (longitudes) (0.0284 cc./m.³ for 172° W. longitude, and 0.0326 cc./m.³ for 158° W. longitude).

2. Significant differences ($P<0.01$) are demonstrated between the means for cruises (0.0366 cc./m.³ for cruise 5 and 0.0244 cc./m.³ for cruise 8).

3. Significant differences ($P<0.02$) are demonstrated among the means for latitudes (these vary from 0.0550

cc./m.³ at the Equator to 0.0150 cc./m.³ at 12° N. latitude).

In this analysis no attempt has been made to isolate the component of variance resulting from time of day at which the hauls were made, and the data are insufficient for a 4-way analysis. In table 14, however, the night samples are rather evenly distributed throughout; e. g., the column for series A, cruise 5, contains 8 night hauls and 1 twilight haul; series A, cruise 8, contains 7 night hauls and 4 twilight hauls; series B, cruise 5, contains 8 night hauls and 2 twilight hauls; and series B, cruise 8, contains 9 night hauls and no twilight hauls. As a consequence we might assume that the day-night variation would tend to cancel out.

To test this assumption we have utilized the values given in table 14, and at the expense of considerable loss of data, we have constructed two 3-way tables, one including only day samples (table 15) and the other only night samples (table 16). An analysis of the day samples reveals quite different results than does an analysis of the night samples or of the combined day and night samples. For the day samples we find no significant differences ($P>0.05$) between series (longitudes), between cruises, or among latitudes. An analysis of the night samples provides the same conclusions as were derived from the combined day and night samples, there being no significant differences ($P>0.05$) between series (longitudes) but highly significant differences ($P<0.01$) between cruises and among latitudes.

TABLE 15.—*Analysis of variance of volumes (cc./m.³) of zooplankton samples taken in day hauls, cruises 5 and 8*

[Latitudes grouped to provide a value for each column; three criteria of classification]

| Latitudes | Series A (172° W. longitude) | | Series B (158° W. longitude) | | Mean |
|---|---|---|---|---|---|
| | Cruise 5 | Cruise 8 | Cruise 5 | Cruise 8 | |
| 15° and 13° N. | 0.0216 | 0.0100 | 0.0158 | 0.0236 | 0.0178 |
| 12° and 11° N. | .0084 | .0097 | .0242 | .0094 | .0129 |
| 10° and 9° N. | .0252 | .0094 | .0225 | .0166 | .0184 |
| 7° and 6° N. | .0204 | .0242 | .0258 | .0477 | .0295 |
| 5° and 4° N. | .0385 | .0367 | .0783 | .0118 | .0413 |
| 3° and 2° N. | .0330 | .0435 | .0658 | .0146 | .0392 |
| 1° N. and 0° | .0556 | .0398 | .0347 | .0220 | .0380 |
| 1° and 2° S. | .0742 | .0109 | .0239 | .0478 | .0392 |
| 3° and 4° S. | .0230 | .0146 | .0239 | .0257 | .0218 |
| Mean | .0333 | .0221 | .0350 | .0244 | |

Least significant difference for latitudinal means=.0233

TABLE 15.—*Analysis of variance of volumes (cc./m.³) of zooplankton samples taken in day hauls, cruises 5 and 8*—Continued

| Source of variation | Degrees of freedom | Sum of squares | Mean square | *F* | *P* |
|---|---|---|---|---|---|
| Main effects: | | | | | |
| Series (S) | 1 | 0.00003481 | 0.00003481 | 0.14 | >0. |
| Cruises (C) | 1 | .00107584 | .00107584 | 4.20 | >0. |
| Latitudes (L) | 8 | .00395996 | .00049500 | 1.93 | >0. |
| First-order interactions: | | | | | |
| S×C | 1 | | | | |
| S×L | 8 | | | | |
| C×L | 8 | .00640043 | .00025602 | | |
| Second-order interaction: | | | | | |
| S×C×L | 8 | | | | |
| Total | 35 | .01147104 | | | |

TABLE 16.—*Analysis of variance of volumes (cc./m.³) zooplankton samples taken in night hauls, cruises and 8*

[Latitudes grouped to provide a value for each column; three criteria classification]

| Latitudes | Series A (172° W. longitude) | | Series B (158° W. longitude) | | Mean |
|---|---|---|---|---|---|
| | Cruise 5 | Cruise 8 | Cruise 5 | Cruise 8 | |
| 13°, 12°, and 11° N. | 0.0374 | 0.0192 | 0.0327 | 0.0310 | 0.0 |
| 10°, 9° N. | .0301 | .0127 | .0531 | .0251 | .0 |
| 8°, 7°, and 6° N. | .0338 | .0139 | .0449 | .0190 | .0 |
| 5°, 4°, and 3° N. | .0410 | .0369 | .0701 | .0255 | .0 |
| 2°, 1° N., and 0° | .0604 | .0548 | .0880 | .0370 | .0 |
| 1°, 2°, and 3° S. | .0664 | .0394 | .0490 | .0558 | .0 |
| Mean | .0448 | .0295 | .0563 | .0322 | |

Least significant difference for latitudinal means=.0

| Source of variation | Degrees of freedom | Sum of squares | Mean square | *F* | |
|---|---|---|---|---|---|
| Main effects: | | | | | |
| Series (S) | 1 | 0.00030246 | 0.00030246 | 2.67 | > |
| Cruises (C) | 1 | .00233248 | .00233248** | 20.62 | < |
| Latitudes (L) | 5 | .00364125 | .00072825** | 6.44 | < |
| First-order interactions: | | | | | |
| S×C | 1 | | | | |
| S×L | 5 | | | | |
| C×L | 5 | .00180978 | .00011311 | | |
| Second-order interaction: | | | | | |
| S×C× L | 5 | | | | |
| Total | 23 | .00808597 | | | |

**Indicates a highly significant ($P<0.01$) mean square value.

Although it has been possible to utilize only portion of the data, the differences between crui and among latitudes may be regarded as real, they have been demonstrated not only in table where the day-night component was present, also in the last analysis, table 16, where the d night component was removed. The evidence t there is less latitudinal variation in the samples than in the night samples introdu a new feature and reveals another possible sou of variation in the data. The cause, or causes this phenomenon, if real, are as yet obscure the authors, but may be related to difference

sic productivity, depth of thermocline, or other ctors of the environment.

In summary, through the foregoing analyses, e have demonstrated that the rich zooplankton tches from near the Equator were significantly eater than those of higher latitudes. We have arned also that there was little difference be- veen the populations along the two longitudes, 2° W. and 158° W., over the range of latitudes mpled. We can state with assurance that there as a distinct difference between the amount of oplankton taken on the two cruises, 5 and 8. he mean for cruise 5 (0.0366 cc./m.[3]) conducted ring the northern summer, was greater than that r cruise 8 (0.0244 cc./m.[3]) conducted during the inter; therefore, a seasonal difference is sug- sted.

## CORRELATIONS WITH ENVIRONMENTAL FACTORS

Some of the hydrographic data obtained on uises of the *Hugh M. Smith* have been pub- shed (Cromwell 1951), and since the Pacific ceanic Fishery Investigations plans to publish e hydrographic data on which these correlations e based, we have not included the data in this per. Variations in certain features of the sur- ce layer as measured on cruise 5, and graphi- lly portrayed in figure 9, may be summarized as llows:

1. Surface inorganic phosphates—high concentra- ions of 0.80 to 0.90 μg at./L. were found in the immedi- ate vicinity of the Equator; values decreased to the orthward and southward to lows of 0.30 to 0.40 μg t./L.

2. Surface temperature—highest water tempera- tures, about 28° C., were recorded for latitudes 6° to 8° N.; temperature decreased gradually to the northward and to the southward, reaching 24° to 25° C.

3. Dissolved oxygen (percent saturation at sur- face)—ranged from 94 percent near the Equator to 102 percent at higher latitudes.

4. Thermocline depth—varied from about 120 feet n the north to over 500 feet in the region of the con- vergence.

Employing cruise-5 data,[6] a method of correla- n analysis (following Snedecor 1946, p. 138) s used to examine the relation between zoo- nkton volumes and these environmental fac- tors. The results, summarized in table 17, point out the following:

1. Statistically significant ($P<0.01$) positive correlations between zooplankton volume and inorganic phosphates at the surface, at the 100-meter depth, and at depths midway to the top of the thermocline;

2. Statistically nonsignificant ($P>0.05$) correlations between zooplankton volume and temperature, whether at the surface, the 100-meter depth, or at depths midway to the top of the thermocline;

3. Statistically significant negative correlations between zooplankton volume and oxygen (as percent saturation) at the surface ($P<0.05$), at the 100-meter depth ($P<0.01$), and at depths midway to the top of the thermocline ($P<0.01$);

4. Statistically significant ($P<0.01$) correlations between zooplankton volume and depth to the top of the thermocline.

In most ecological investigations it has been found that all factors examined are more or less interrelated, variations in one factor having an influence, either direct or indirect, on all the other factors (Riley 1939a). The multiple-regression method makes corrections for such interactions, and in a three-variate analysis tests the relation of a pair of variates irrespective of the influence of a third.

TABLE 17.—*Correlations of zooplankton abundance, as wet volumes, and certain environmental factors, cruise 5*

| Variates | | Degrees of freedom | Correlation coefficient ($r$) | $P$ |
|---|---|---|---|---|
| $X_1$ | $X_2$ | | | |
| Inorganic phosphates (μg at./L.) at surface. | Zooplankton wet volumes. | 41 | 0.655** | <0.01 |
| Inorganic phosphates (μg at./L.) at 100 m. depth. | do | 41 | 0.580** | <0.01 |
| Inorganic phosphates (μg at./L.) at depths midway to thermocline. | do | 41 | 0.730** | <0.01 |
| Temperature (° C.) at surface | do | 49 | 0.008 | >0.05 |
| Temperature (° C.) at 100 m. depth. | do | 49 | 0.227 | >0.05 |
| Temperature (° C.) at depths midway to thermocline. | do | 49 | −0.065 | >0.05 |
| Oxygen (percent saturation) at surface. | do | 49 | −0.291* | <0.05 |
| Oxygen (percent saturation) at 100 m. depth. | do | 49 | −0.834** | <0.01 |
| Oxygen (percent saturation) at depths midway to thermocline. | do | 49 | −0.489** | <0.01 |
| Thermocline depth (feet) | do | 49 | 0.566** | <0.01 |

** Indicates a highly significant correlation.
* Indicates a significant correlation.

In the correlation analyses a positive relation between zooplankton volumes and temperature at the 100-meter depth was indicated (table 17), but, being below the 0.05 level of probability, it was not considered statistically significant. It seemed of interest to examine the relation of zooplankton

[6] Since different methods of towing and nets of different mesh were used on cruise 2, and since the hydrographic data of ise 8 are not as yet entirely processed, we have used cruise-5 a for the correlation analyses.

and temperature at the 100-meter level independent of variation with inorganic phosphate. A multiple regression analysis (following Snedecor 1946, pp. 340–373) was therefore carried out employing cruise-5 data, treating zooplankton volumes (cc./m.$^3$) as the $Y$ variate, inorganic phosphates ($\mu$g at./L) at 100 meters as the $X_1$ variate, and temperatures (°C.) at 100 meters as the $X_2$ variate.

The following multiple regression equation was obtained:

$$Y=0.0484X_1+0.0023X_2-0.044969.$$

The first partial regression coefficient ($b_{Y1\cdot2}$= 0.0484; $t$=5.232, $P<0.001$, at 40 degrees of freedom) and the first partial correlation coefficient ($r_{Y1\cdot2}$=0.638; $P<0.01$) are highly significant, showing that there is a positive relation between zooplankton volumes and inorganic phosphates, independent of the variation with temperature, at the 100-meter level. This conclusion was expected from the previous correlation analysis. The second partial regression coefficient ($b_{Y2\cdot1}$=0.0023; $t$=2.987, $P<0.01$, at 40 degrees of freedom) and the second partial correlation coefficient ($r_{Y2\cdot1}$= 0.423; $P<0.01$) are also highly significant, showing that there is also a positive relation between zooplankton volumes and temperature, independent of variation with inorganic phosphate, at the 100-meter level. This conclusion is of interest since it indicates a much higher degree of covariation between zooplankton and temperature, than was revealed by the previous correlation analysis.

We have shown that the abundance of zooplankton in the central Pacific is correlated with inorganic phosphates, temperature, dissolved oxygen, and thermocline depth. We do not believe, though, that the abundance of zooplankton is in any way limited by the conditions of temperature, oxygen, or thermocline depth prevailing throughout the region. The relationships between zooplankton and these three environmental factors, as indicated by the statistical correlations, are most likely independent variations due to common causes. Zooplankton is more directly linked with phosphate, however, through the phosphate→ phytoplankton→zooplankton relation. Therefore, the causal agent for these several variations examined is directly or indirectly the Equatorial Current system, principally the divergence at the Equator which brings to the surface water that is relatively low in temperature, low in oxygen co tent, but high in chemical nutrients, and the co vergence to the northward which results in deepening of the thermocline and a possible co centration of the plankton.

Other investigators have found essentially sim lar correlations between plankton and these e vironmental factors. Marshall and Orr (192 observed in the Clyde Sea area that where anim life was rich, phosphates were high, but dissolv oxygen and pH were low. Hardy and Gunth (1935) found in the Pacific that numbers of zo plankton were positively correlated with ph phate values. Jesperson (1935) states that the is a direct relation between quantities of nitra and phosphate and macroplankton. Leav (1938) found in the Atlantic Basin a correlati between temperature, salinity, and density, a the vertical distribution of zooplankton; a neg tive correlation was found between oxygen a zooplankton. Graham (1941) reports that diff ences in productivity in the Pacific, as measured plankton dry weight, are correlated with the c centration of phosphate.

## COMPARISON WITH ZOOPLANKTON ABUNDAN OF OTHER REGIONS

To facilitate the comparison of our data w that of other regions, we have calculated avera numbers and volumes (table 11) which are rep sentative of the quantity of zooplankton in diff ent latitudinal zones of the central Pacific. I a difficult task, however, to reduce to compara terms plankton data which have been obtained different investigators, using different methods different seasons, and with results expressed in c ferent units. In surveying a considerable fract of the great bulk of available literature on qu titative plankton sampling, we found only a f reports that have enough in common with our o work to permit a comparison of the results.

In Graham's report (1941) on plankton lected by the *Carnegie* along a series of stati extending from San Francisco to Samoa, he sta "In the open Pacific Ocean from September November 1929, there was a greater productio total plankton in the tropics between latitu 20° N. and 11° S. than between latitudes 20° 34° N." The richest collection of this section 10) was taken at 13° N. latitude, but was con ered atypical since it consisted largely of sa




IGURE 10.—Station positions and zooplankton abundance (as dry weight of sample in milligrams) found on the last cruise of the *Carnegie* in 1929 (data from Graham 1941).

he sample ranking second in dry weight came om 5°30′ N. latitude and the third-ranking sam- le from 2° S. latitude. These results, while based n a different method of analysis, are in general greement with our own observations.

The Japanese have carried on extensive plank- n investigations in waters adjacent to the Palau lands and in the area between the Palaus and ew Guinea. Results from the studies of aneda (1942) and Tokioka (1942) are illus- ated in figure 11. Although these data are ther limited in scope, the samples of Tokioka show quite definitely the influence of enrichment near the Equator.

These surveys and the work of the *Carnegie* are the only north-south plankton studies that we are aware of in the equatorial Pacific, other than our own. Unfortunately, the absolute values obtained in them cannot be compared with our results, but the generally similar variations with latitude are of interest.

While crossing the Pacific from Panama to the Indo-Pacific region, the *Dana* expedition made plankton tows at frequent intervals. Jesperson




FIGURE 11.—Plankton abundance in the western Pacific: Results given are settlement volumes of total (?) plank obtained by vertical hauls at 50 to 0 meters along 134° E. longitude. A. As reported by Tokioka (1942) three cruises, December 1939, March 1940, and May 1940 (averaged by the senior author). B. As reported Haneda (1942) for one cruise, December 1939.

(1935), summarizes the results as follows (the volumes expressed are catches per 1-hour haul using 1½-meter stramin nets with 50 and 100 meters of wire out):

> A notable fact here is, that we find specially large volumes in the eastern part of the Pacific, from the Bay of Panama to the Marquesas Is., with a distinct maximum (1,125 cc.) at St. 2,558, which lies west of the Galapagos. On the stretch from the Marquesas to Tahiti we have decreasing quantities of plankton (ca. 150–200 cc.), whilst on the sections from Tahiti—Cook Island (Rarotonga)—Samoa—Fiji the quantities are very small, less than 100 cc. We thus have extremely little macroplankton in this area of the central part of the Pacific, yet a series of stations just north of Samoa yielded somewhat larger quantities (ca. 100–260 cc.). From Fiji to New Caledonia also the quantity of macroplankton increases (ca. 125–210 cc.) and a further increase is shown in the section from New Caledonia down towards Kermadec Islands.

For comparison, in the North Atlantic the expedition obtained volumes ranging from 90 to 7,250 cc. per hour of hauling.

The very extensive plankton data presently b ing collected by the California Cooperative S dine Research Program (California, Progress R port 1950, and unpublished data) are quite co parable to our own in most respects. Similar n have been used by both investigations, and the sults are expressed in similar units. Whereas t Sardine Research Program employed an oblic tow to a depth of 70 meters for the collections ported here, our oblique tow descended to 2 meters. This difference in sampling method pr ably had no great influence on the difference in sults obtained. If we consider 22 of their farth offshore stations, located between 25° N. and 3 N. latitude and visited in September 1950, we fi that the average for the group was 0.057 cc./m which is approximately equivalent to our avera (table 11) for the South Equatorial Current n the Equator but 3 times the average for the No Equatorial Current. Values for regions of u welling close in to the California coast were high as 14.595 cc./m.³ in February 1950. This

any times our maximum value of 0.1025 cc./m.³ he average of 0.208 cc./m.³ for the entire West oast between 25° N. and 47° N. in April and May 949, is just about 10 times our mean for the cen- al equatorial Pacific.

Other regions of sparse plankton are found in ie Gulf Stream and in the Sargasso Sea of the tlantic Ocean. These areas apparently have uch poorer plankton populations than Atlantic astal and continental shelf waters (Clarke 1940; iley 1939b; Riley and Gorgy 1948).

Table 18 is a summary of representative plankton values for various areas of the Atlantic and Pacific Oceans. Part of the variation in these values may be due to differences in mesh size among the nets used by different investigators. The averages for the central Pacific are the poorest of the lot; in fact they are so low that one is inclined to speculate as to how the pelagic fish populations, particularly the relatively large population of tunas, are supported.

TABLE 18.—*Comparison of plankton abundance in various areas of the Atlantic and Pacific Oceans*

[Values given in wet volume, estimated number, or wet weight]

| Area | Plankton values | Mesh aperture of net used | Reference |
|---|---|---|---|
| CIFIC AREA: | | | |
| Samoa to the Marshalls via the Equator | 0.4 to 0.95 cc./m.³ | 0.33 mm | Kramer (1906). |
| Java Sea | Avg.=0.83 cc./m.³ | 0.33 mm | Delsman (1939). |
| Open sea, Palau Is. | 110 to 530, no./m.³ | 0.37 mm | Motoda (1940). |
| Bikini, just outside reef | 26.23 to 62.20 no./m.³ | | Johnson (1949). |
| West coast, U. S. (offshore stations) | 0.016 to 0.129 cc./m.³<br>Avg.=0.057 cc./m.³ | 0.65 mm | California prog. rept. (1950). |
| Central Pacific | 5 to 109, no./m.³ Avg.=35<br>0.002 to 0.102 cc./m.³<br>Avg.=0.027 cc./m.³ | 0.65 mm | This report. |
| LANTIC AREA: | | | |
| Baltic Sea | 1.9 to 11.0 cc./m.³ | 0.33 mm | Kramer (1906). |
| Gulf of Maine (10 to 100 m. depth) | 0.12 to 4.30 cc./m.³ | 1.25 mm., front; 0.8 mm., middle and rear | Bigelow (1926). |
| North Atlantic (coastal water) | Avg.=0.5 to 0.8 cc./m.³ | 29.38 meshes/in., front; 48.54 meshes/in., rear | Bigelow and Sears (1939). |
| Florida Strait | 0.02 cc./m.³ | | |
| Gulf Stream, off Florida | 0.05 cc./m.³ | | |
| Gulf Stream, off Georgia | 0.07 cc./m.³ | 0.158 mm | Riley (1939b). |
| North Atlantic continental slope | 4.3 cc./m.³ | | |
| North Atlantic, coastal | 8.1 cc./m.³ | | |
| North Atlantic, coastal | Avg.=0.54 cc./m.³<br>Max.=15.5 cc./m.³ | 10 strands/cm | Clark (1940). |
| North Atlantic, offshore | Avg.=0.40 cc./m.³<br>Max.=3.5 cc./m.³ | | |
| Sargasso Sea | Avg.=0.045 gm./m.³ | 0.158 mm | Riley and Gorgy (1948). |
| Gulf Stream | 1 sta.=0.137 gm./m.³ | | |
| Slope water | 2 sta.=0.14 and 1.6 gm./m.³ | | |

## PRODUCTIVITY

The practical application of most plankton re- arch is to provide data for estimating and com- aring the productivity, or available food, in arious areas of the sea. It has been strongly em- hasized in more recent plankton literature that e "standing crop" does not give a true measure f the rate of production. Harvey (1934) and arvey et al. (1935) have shown that the size of e standing crop of phytoplankton is greatly af- cted by the grazing of animal herbivores and erefore at any one time is merely a momentary lance between the processes of production and nsumption. In the tropics, steady grazing by redators may keep the zooplankton at a lower vel of abundance than in higher latitudes where asonal features of the environment allow the ankton to "pulse" or bloom and thus increase uch faster than the predators. The apparently low standing crop may be considerably counterbalanced by a high rate of turnover and nearly uniform production throughout the year.

It is generally assumed that in water masses where the annual plant production is great the density of the animal population will also be great. This assumption is roughly borne out by general observations (Harvey 1945). Delsman (1939) has stated, "Where no rich plankton can develop, no rich macrofauna, no abundant fish population can either be expected." In the same vein, "The dependence of various elements of the food chain on a preceding one, conditions the distribution of the larger forms" (Hesse, Allee, and Schmidt 1951). Also, it is reasonable to believe that the zooplankton population will be the maximum that the plant crop can support. Local situations may not conform to this generality, but when large areas are considered, there is usually found a direct

relation between concentrations of phytoplankton and zooplankton (Riley and Bumpus 1946). In conducting a fish-farming experiment, Raymont (1947) found that in both enclosed and unenclosed small sea areas the addition of a nitrate and phosphate fertilizer stimulated phytoplankton growth, which in turn maintained a high density of zooplankton which promoted a rapid growth of flatfish.

From tuna-catch records and observations on occurrence of surface schools, the Pacific Oceanic Fishery Investigations is accumulating evidence on the distribution of tunas which indicates quite definitely that areas of the greatest zooplankton abundance in the central Pacific are also areas of greatest tuna abundance.

We are fully aware that the data we have presented on the variation of zooplankton abundance with latitude does not in any way reflect the "rate of turnover," the most difficult element to determine in estimates of productivity. In this area of the tropical Pacific, with temperatures very uniform in time and space and zooplankton very uniform in composition, the rate of turnover should not be a disturbing feature in the comparison of the several parts of our area, i. e., for our data standing crop should be proportional to productivity.

## SUMMARY AND CONCLUSIONS

1. This report presents the results of 210 quantitative zooplankton collections made in the central Pacific in 1950 and 1951, between 27° N. and 14° S. latitude, and 155° and 175° W. longitude.

2. Most of the collections were obtained by oblique hauls to 200 meters' depth, employing 1-meter nets of 30xxx grit gauze with aperture widths of 0.65 mm.

3. A method of sampling was developed that harmonized with hydrographic and fishing operations, required little of ship's time, and involved no particularly elaborate treatment of samples in the laboratory.

4. The zooplankton taken by our collecting method was composed on the average by number, of 57 percent Copepoda, with the other chief components ranking as follows: Chaetognatha, 12 percent; Tunicata, 6 percent; Euphausiacea, 5 percent; Siphonophora, 4 percent; and Foraminifera, 4 percent.

5. The composition of the collections was r markably uniform when longitudes and cruis were compared.

6. Despite the use of oblique tows to 200 mete depth, we found significant differences betwe day hauls and night hauls.

7. Since distinctly larger catches were obtain on the "summer" cruise (cruise 5) than on t "winter" cruise (cruise 8), a seasonal difference zooplankton abundance is indicated.

8. Within the range of latitudes sampled, t greatest abundance, both by number and volum of zooplankton occurred in the region of the Equ tor; sometimes the greatest concentrations we found north of the Equator, when related to convergence, and to the south when no mark convergence existed.

9. The abundance of zooplankton is correlat with such chemical and physical environment factors as inorganic phosphate, water temper ture, dissolved oxygen, and thermocline dept which are influenced by the upwelling resulti from the equatorial divergence.

10. Upwelling along the Equator replenish the supply of nutrients in the euphotic zone, th providing a favorable environment for the grow of phytoplankton. Since animal life fluctuat with its food supply, conditions in this regi are favorable for the development of a zooplan ton population.

11. While our observations on the standi crop of zooplankton do not give a measure of t rate of production in its strict sense, we belie that they do provide a useful index to the relati productivity of different areas of the centr Pacific.

## LITERATURE CITED

AGASSIZ, ALEXANDER, and A. G. MAYER.
1902. Report on the scientific results of the expediti to the Tropical Pacific. III. Medusae. Mus. Co Zool., Mem. 26 (3) : 139–176.

ARRHENIUS, GUSTAF.
1950. Late cenozoic climatic changes as recorded the Equatorial Current System. Tellus 2 (2 83–88.

BEEBE, WILLIAM.
1926. The Arcturus Adventure. xix, 439 p. N York: G. P. Putnam's Sons.
1936. Food of the Bermuda and West Indian tur of the genera *Parathunnus* and *Neothunnus*. Z logica 21 (15) : 196–205.

ELOW, H. B.
926. Plankton of the offshore waters of the Gulf of Maine. Bull. U. S. Bur. Fish. (1924), 40 (2): 1–509.

ELOW, H. B., and MARY SEARS.
939. Studies of the waters of the Continental Shelf, Cape Cod to Chesapeake Bay. III. A volumetric study of the zooplankton. Mus. Comp. Zool., Mem. 54 (4): 183–378.

OOKS, S. C.
934. Ocean currents and the migration of pelagic birds. Condor 36 (5): 185–190.

LIFORNIA COOPERATIVE SARDINE RESEARCH PROGRAM.
950. Progress Report, 1950. State of California, Dept. Nat. Resources, Mar. Research Comm. 54 p.

ARKE, GEORGE L.
940. Comparative richness of zooplankton in coastal and offshore areas of the Atlantic. Biol. Bull. 78 (2): 226–255.
946. Dynamics of production in a marine area. Ecol. Monogr. 16 (4): 321–335.

EMENS, W. A., and G. V. WILBY.
946. Fishes of the Pacific Coast of Canada. Fish. Res. Bd. Canada, Bull. No. 68, 368 p.

OMWELL, TOWNSEND.
951. Mid-Pacific oceanography, January through March, 1950. U. S. Fish and Wildlife Serv., Spec. Sci. Rept.: Fisheries No. 54, 79 p. 17 figs.

LSMAN, H. C.
939. Preliminary plankton investigations in the Java Sea. Treubia 17 (2): 139–181.

RDINER, A. C.
931. The validity of single vertical hauls in the study of the distribution of the plankton. Jour. Mar. Biol. Assoc., (n. s.), 17 (2): 449–472.

AHAM, HERBERT W.
941. Plankton production in relation to character of water in the open Pacific. Jour. Mar. Res. 4 (3): 189–197.

ECKEL, ERNST.
890. Plankton studies. Jenaische Zeitschrift, XXV. Translated by G. W. Field as Appendix 6 to the Report of the Commissioners for 1889 to 1891, U. S. Comm. Fish and Fisheries, Washington, 1893, pp. 565–641.

NEDA, RYŌKA.
942. Ocean currents and plankton of waters adjacent to the Palau Islands. South Sea Sci. [Kagaku Nanyō] 5 (1): 78–84. (Translated from the Japanese by Lawrence B. McNeill, Jr.)

RDY, A. C.
936. Observations on the uneven distribution of oceanic plankton. Discovery Reports 11: 511–538.

RDY, A. C., and E. R. GUNTHER.
935. The plankton of the South Georgia Whaling Grounds and adjacent waters, 1926–27. Discovery Reports 11: 1–456.

HARVEY, H. W.
1934. Annual variation of planktonic vegetation. Jour. Mar. Biol. Assoc., (n. s.) 19 (2): 775–792.
1945. Recent advances in the chemistry and biology of sea water. vii, 164 p. London: Cambridge Univ. Press.

HARVEY, H. W., L. H. N. COOPER, M. V. LEBOUR, and F. S. RUSSEL.
1935. Plankton production and its control. Jour. Mar. Biol. Assoc., (n. s.) 20 (2): 407–441.

HERDMAN, W. A.
1923. The founders of oceanography and their work. xii, 340 p. London: Edward Arnold and Co.

HESSE, RICHARD, W. C. ALLEE, and KARL P. SCHMIDT.
1951. Ecological animal geography. 2nd ed., xiii, 715 p. New York: John Wiley and Sons, Inc.

IMAMURA, YUTAKA.
1949. The Japanese skipjack fishery. Suisan Kōza, vol. 6, Fishing Section, pp. 17–94. Published by Dai Nippon Suisan Kai [Japanese Fisheries. Assoc.], Tokyo. (Translated from the Japanese by W. G. Van Campen. United States Fish and Wildlife Service, Spec. Sci. Rept.: Fisheries No. 49, 67 p., 1951.)

IVLEV, V. S.
1945. The biological productivity of waters. Uspekhi Sovremennoi Biologii (Advances in Modern Biology), 19 (1): 98–120. (Translated from the Russian by W. E. Ricker.)

JESPERSON, P.
1935. Quantitative investigations on the distribution of macroplankton in different ocean regions. Rept. "Dana" Exped. 2 (7): 1–44.

JOHNSON, L. P. V.
1950. An introduction to applied biometrics. vi, 165 p. Minneapolis: Burgess Publishing Co.

JOHNSON, M. W.
1949. Zooplankton as an index of water exchange between Bikini Lagoon and the open sea. Trans. Amer. Geophys. Un. 30 (2): 238–244.

KENDALL, MAURICE G.
1948. The advanced theory of statistics. Vol. II, 2nd ed., vii, 521 p. London: Charles Griffin and Co., Ltd.

KISHINOUYE, KAMAKICHI.
1917. The food of tunas. Suisan Gakkai Hō 2 (1): 106–108. (Translated from the Japanese by W. G. Van Campen. Translation No. 29, Pacific Oceanic Fishery Investigations, United States Fish and Wildlife Service, Washington, D. C.)
1924. Observations on the skipjack fishing grounds. Suisan Gakkai Hō 4 (2): 87–92. (Translated from the Japanese by W. G. Van Campen. Translation No. 21, Pacific Oceanic Fishery Investigations, United States Fish and Wildlife Service, Washington, D. C.)

KRAMER, A.
1906. Ergebnisse meiner Korallenriff- ünd Planktonstudien. [Cited by Delsman, H. C. 1939. Preliminary plankton investigations in the Java Sea. Treubia 17 (2): 139–181.]

LEAVITT, B.
1938. The quantitative vertical distribution of macrozooplankton in the Atlantic Ocean basin. Biol. Bull. 74 (3) : 376–394.

MARSHALL, S. M.
1933. The production of microplankton in the Great Barrier Reef Region. Gr. Barrier Reef Exped. 1928–29, Sci. Rept. 2 (5) : 111–157.

MARSHALL, S. M., and A. P. ORR.
1927. The relation of the plankton to some chemical and physical factors in the Clyde Sea area. Jour. Mar. Biol. Assoc. (n. s.), 14 (4) : 837–868.

MATSUYA, KENZŌ.
1937. Some hydrographical studies of the water of Iwayama Bay in the South Sea Islands. Palao Trop. Biol. Sta. Studies 1 (1) : 95–135.

MOTODA, S.
1940. Comparison of the conditions of water in the bay, lagoon, and open sea in Palau. Palao Trop. Biol. Sta. Studies 2 (1) : 41–48.

NAKAI, Z.
1942. The chemical composition, volume, weight, and size of the important marine plankton. Jour. Oceanogr. Soc. Japan 1 : 45–55, 146 (English abstract).

RAYMONT, J. E. S.
1947. A fish-farming experiment in Scottish sea lochs. Jour. Mar. Res. 6 (3) : 219–222.

REINTJES, J. W., and J. E. KING.
1953. Food of yellowfin tuna in the Central Pacific. United States Fish and Wildlife Service, Fishery Bull. 54 (81) : 91–110.

REVELLE, R.
1944. Marine bottom samples collected in the Pacific Ocean by the *Carnegie* on its seventh cruise. Carnegie Inst. Wash., Pub. 556 : 1–180.

RILEY, GORDON A.
1939a. Correlations in aquatic ecology with an example of their application to problems of plankton productivity. Jour. Mar. Res. 2 (1) : 56–73.
1939b. Plankton studies. II. The western North Atlantic, May–June, 1939. Jour. Mar Res. 2 (2) : 145–162.

RILEY, G. A., and D. F. BUMPUS.
1946. Phytoplankton-zooplankton relationships on Georges Bank. Jour. Mar. Res. 6 (1) : 33–47.

RILEY, GORDON A., and SAMY GORGY.
1948. Quantitative studies of summer plankton populations of the western North Atlantic. Jour. Mar. Res. 7 (2) : 100–121.

RUSSEL, F. S.
1934. The zooplankton. III. A comparison of t abundance of zooplankton in the Barrier Reef Lago with that of some regions in northern Europe waters. Gr. Barrier Reef Exped. 1928–29, Sci. Re 2 (6) : 176–185.

RUSSEL, F. S., and J. S. COLMAN.
1931. The zooplankton. I. Gear, methods, and s tion lists. Gr. Barrier Reef Exped. 1928–29, S Rept. 2 (2) : 5–35.
1934. The zooplankton. II. The composition of t zooplankton of the Great Barrier Reef Lagoon. Barrier Reef Exped. 1928–29, Sci. Rept. 2 (6) : 1 201.
1935. The zooplankton. IV. The occurrence and s sonal distribution of the Tunicata, Mollusca, a Coelenterata (Siphonophora). Gr. Barrier R Exped. 1928–29, Sci. Rept. 2 (7) : 203–276.

SARGENT, M. C., and T. S. AUSTIN.
1949. Organic productivity of an atoll. Trans. Am Geophys. Un. 30 (2) : 245–249.

SEARS, MARY.
1950. Notes on siphonophores. I. Siphonopho from the Marshall Is. Jour. Mar. Res. 9 (1) : 1–

SNEDECOR, GEORGE W.
1946. Statistical methods. 4th ed., xvi, 485 p. Am Iowa State College Press.

STEEMANN NIELSON, E.
1937. On the relation between quantities of phy plankton and zooplankton in the sea. Jour. Co Int. Expl. Mer. 12 (2) : 147–154.

SUYEHIRO, YASUO.
1942. A study on the digestive system and feed habits of fish (in English). Japanese Jour. Z 10 (1) : 1–303.

SVERDRUP, H. U., M. W. JOHNSON, and R. H. FLEMING
1942. The oceans, their physics, chemistry, and gene biology. x, 1087 p. New York : Prentice-Hall, Inc

TOKIOKA, TAKASHI.
1942. Plankton abundance in Iwayama Bay a waters surrounding the Palau Islands. South Sci. [Kagaku Nanyō] 5 (1) : 44–55. (Transla from the Japanese by Lawrence B. McNeill, Jr.)

WILSON, CHARLES B.
1942. Copepods of the plankton gathered during last cruise of the *Carnegie*. Carnegie Inst. Wa ington, Pub. 536 : 1–237.

WINSOR, C. P., and G. L. CLARKE.
1940. A statistical study of variation in the catch plankton nets. Jour. Mar. Res. 3 (1) : 1–34.